

## ALGORITHMS FOR L-CONVEX FUNCTION MINIMIZATION: CONNECTION BETWEEN DISCRETE CONVEX ANALYSIS AND OTHER RESEARCH FIELDS

Akiyoshi Shioura  
*Tokyo Institute of Technology*

(Received July 19, 2016; Revised October 24, 2016)

*Abstract* L-convexity is a concept of discrete convexity for functions defined on the integer lattice points, and plays a central role in the framework of discrete convex analysis. In this paper, we review recent development of algorithms for L-convex function minimization. We first point out the close connection between discrete convex analysis and various research fields such as discrete optimization, auction theory, and computer vision by showing that algorithms proposed independently in these research fields can be regarded as minimization algorithms applied to specific L-convex functions. Therefore, we can provide a unified approach to analyze the algorithms appearing in various research fields through the concept of L-convex function. We then present the recent results on theoretical bounds of the number of iterations required by some minimization algorithms, where precise bounds are given in terms of distance between the initial solution and the minimizer found by the algorithms. From these results we see that the algorithms output the “nearest” minimizer to the initial solution, and that the trajectories of solutions generated by the algorithms are “shortest paths” from the initial solution to the found minimizer. Finally, we consider an application of the results to iterative auctions in auction theory. We point out that the essence of the iterative auctions proposed by Ausubel (2006) lies in L-convexity. We also present new iterative auctions by Murota–Shioura–Yang (2016), which are based on the understanding of existing iterative auctions from the viewpoint of discrete convex analysis.

**Keywords:** Discrete optimization, discrete convex function, steepest descent algorithm, analysis of algorithm, iterative auction

### 1. Introduction

L-convexity and  $L^{\natural}$ -convexity are concepts of discrete convexity for functions defined on the integer lattice points. The concept of L-convexity is introduced by Murota [31]. As a variant of L-convexity, the concept of  $L^{\natural}$ -convexity (read “L-natural-convexity”) is introduced by Fujishige–Murota [15]. The concepts of  $L/L^{\natural}$ -convexity are later extended to functions in continuous variables [39].  $L/L^{\natural}$ -convexity plays a central role in discrete convex analysis initiated by Murota [31, 32], which is a theoretical framework for efficiently solvable discrete optimization problems.

In this paper, we consider the minimization of  $L/L^{\natural}$ -convex functions. Such a problem can be found in various research fields. For example, discrete optimization problems such as the shortest path problem and the dual of the minimum cost flow problem can be seen as special cases of L-convex function minimization [16, 43, 44]. Moreover, L-convex function minimization can also be found in other research fields such as computer vision (see Section 3.3.2), auction theory (see Section 6), inventory theory [24, 53], etc. (see also [32, 33, 37]). This fact indicates that discrete convex analysis is closely connected with various research fields through L-convex function minimization.

It is known that a global minimizer of an L-convex function can be characterized by a local minimality (see Section 2.3). Therefore, minimization of an L-convex function can be solved by steepest descent algorithms [32, 33]. Such algorithms can be found in other research fields; indeed, optimization algorithms proposed independently in different research fields such as discrete optimization, computer vision, and auction theory coincide with some variants of the steepest descent algorithms applied to special L-convex functions (see Section 3.3). Hence, we can provide a unified viewpoint to algorithms appearing in various research fields through the analysis of steepest descent algorithms for L-convex functions.

In this paper, we review the results in [16, 44] concerning the analysis of steepest descent algorithms for L-convex function minimization. In particular, we present theoretical bounds on the number of iterations required by the algorithms, where the precise bounds are given in terms of distance between the initial solution and the set of minimizers. This fact implies that the algorithms output the “nearest” minimizer to the initial solution, and that the trajectories of solutions generated by the algorithms are “shortest paths” from the initial solution to the found minimizer.

The results on the analysis of L-convex function minimization algorithms are interesting on their own right and also useful in other research fields. For example, it is a common approach in computer vision to process some tasks for images by solving the minimization of a certain energy function, which can be seen as a special case of L-convex function minimization. It is known that when we solve such a minimization problem, simple algorithms similar to a steepest descent algorithm run fast in practice. Since the running time of the steepest descent algorithm is heavily dependent on the number of iterations of the algorithm, we can provide a theoretical guarantee for the running time through a bound on the number of iterations.

As an interesting application of the results on L-convex function minimization algorithms, we consider iterative auctions in auction theory [45, 46]. An iterative auction is an algorithm (protocol, more precisely) for computing equilibrium prices of goods by iteratively changing prices with the aid of the bidders’ reported information. Recently, Murota–Shioura–Yang [45, 46] pointed out the connection between iterative auctions and discrete convex analysis by showing that iterative auctions for certain auction models can be regarded as minimization algorithms for specific  $L^{\natural}$ -convex functions. (see also Drexl–Kleiner [13]). We explain this connection, and show that how the results in the previous sections can be applied to iterative auctions in Section 6.

The organization of this paper is as follows. In Section 2 we describe the definition of L-convex function and present fundamental properties of minimizers. In Section 3 various minimization algorithms for L-convex functions are presented. In Section 4, we present theoretical results on the behavior of minimization algorithms for L-convex functions and provide precise bounds for the number of iterations. In Section 5 we consider minimization algorithms for L-convex functions in continuous variables, and observe that some theoretical results for the case of discrete variables extend to the case of continuous variables. Finally in Section 6, we explain an application to iterative auctions.

This paper is based on the manuscript [51] (in Japanese) that appeared in the proceedings of the 27th RAMP Symposium of Operations Research Society of Japan.

## 2. L-convex and $L^{\natural}$ -convex Functions

In this section we present the definitions and examples of L-convex and  $L^{\natural}$ -convex functions.

## 2.1. Definitions

Let  $n$  be a positive integer and denote  $N = \{1, 2, \dots, n\}$ . We denote by  $\mathbb{Z}_+$  and by  $\mathbb{R}_+$  the sets of non-negative integers and non-negative reals, respectively.

A function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is called an *L-convex function* [31] if it satisfies the following two properties:

$$\text{[Submodularity]} \quad g(p) + g(q) \geq g(p \vee q) + g(p \wedge q) \quad (\forall p, q \in \text{dom } g), \quad (2.1)$$

**[Linearity in Direction of  $\mathbf{1}$ ]**

$$\exists r \in \mathbb{R} : g(p + \alpha \mathbf{1}) = g(p) + \alpha r \quad (\forall p \in \mathbb{Z}^n, \forall \alpha \in \mathbb{Z}), \quad (2.2)$$

where  $\text{dom } g = \{p \in \mathbb{Z}^n \mid g(p) < +\infty\}$ ,  $\mathbf{1} = (1, 1, \dots, 1)$ , and  $p \vee q$ ,  $p \wedge q$  denote the vectors obtained by the component-wise maximum and minimum of  $p$  and  $q$ , i.e.,

$$(p \vee q)(i) = \max(p(i), q(i)), \quad (p \wedge q)(i) = \min(p(i), q(i)) \quad (i \in N).$$

Due to the linearity in the direction of  $\mathbf{1}$  (i.e., (2.2)), the essential information of an L-convex function  $g$  is not lost by the restriction of  $g$  to an arbitrarily chosen coordinate hyperplane. That is, for an L-convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n$  variables, the function  $\check{g} : \mathbb{Z}^{n-1} \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n - 1$  variables given by

$$\check{g}(q(1), \dots, q(n-1)) = g(q(1), \dots, q(n-1), 0)$$

has essentially the same information as  $g$ . We call such  $\check{g}$  an *L<sup>1</sup>-convex function* [15]. Hence, a function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is L<sup>1</sup>-convex function if the function  $\tilde{g} : \mathbb{Z}^{n+1} \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n + 1$  variables given by

$$\tilde{g}(q(1), \dots, q(n), q(n+1)) = g(q(1) - q(n+1), \dots, q(n) - q(n+1)) \quad (q \in \mathbb{Z}^{n+1}) \quad (2.3)$$

is an L-convex function. Note that the function  $\tilde{g}$  given by (2.3) always satisfies the property (2.2). Hence,  $\tilde{g}$  is an L-convex function if and only if  $\tilde{g}$  satisfies the submodularity (2.1). That is, L<sup>1</sup>-convexity of a function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is equivalent to the submodularity of  $\tilde{g}$ .

It is known that L<sup>1</sup>-convexity is equivalent to the following property (see Figure 1):

**[Translation Submodularity]**

$$g(p) + g(q) \geq g((p - \alpha \mathbf{1}) \vee q) + g(p \wedge (q + \alpha \mathbf{1})) \quad (\forall p, q \in \text{dom } g, \forall \alpha \in \mathbb{Z}_+). \quad (2.4)$$

L<sup>1</sup>-convexity is also equivalent to a discrete version of mid-point convexity:

**[Discrete Mid-point Convexity]**

$$g(p) + g(q) \geq g\left(\left\lceil \frac{p+q}{2} \right\rceil\right) + g\left(\left\lfloor \frac{p+q}{2} \right\rfloor\right) \quad (\forall p, q \in \text{dom } g), \quad (2.5)$$

where  $\lceil \frac{p+q}{2} \rceil$  and  $\lfloor \frac{p+q}{2} \rfloor$  denote the integral vectors obtained by component-wise rounding up and rounding down of the vector  $\frac{p+q}{2}$ , respectively.

From the discussion above, we obtain the following equivalence.

**Theorem 2.1** ([15]). *For a function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  the following three properties are equivalent:*

- (a) *L<sup>1</sup>-convexity, i.e., L-convexity of  $\tilde{g}$  in (2.3),*
- (b) *translation submodularity (2.4),*
- (c) *discrete mid-point convexity (2.5).*

□




Figure 1: Vectors  $(p - \alpha \mathbf{1}) \vee q$  and  $p \wedge (q + \alpha \mathbf{1})$  in the translation submodularity (2.4): (Left) Case of  $p \geq q$ , (Right) Case of  $p(1) > q(1)$  and  $p(2) < q(2)$ . Trajectories of the two vectors when  $\alpha$  is increased from 0 is drawn by dashed arrows. Note that if  $p \leq q$ , then  $(p - \alpha \mathbf{1}) \vee q = q$  and  $p \wedge (q + \alpha \mathbf{1}) = p$  for every  $\alpha \geq 0$ .

By its definition, the concept of  $L^{\sharp}$ -convex function is essentially equivalent to the concept of L-convex function. On the other hand, L-convex function can be seen as a special case of  $L^{\sharp}$ -convex function. Indeed, translation submodularity of an L-convex function can be shown as follows, where  $p, q \in \text{dom } g$  and  $\alpha \in \mathbb{Z}_+$ :

$$\begin{aligned} g(p) + g(q) &= g(p - \alpha \mathbf{1}) + \alpha r + g(q) \quad (\text{by (2.2)}) \\ &\geq g((p - \alpha \mathbf{1}) \vee q) + g((p - \alpha \mathbf{1}) \wedge q) + \alpha r \quad (\text{by (2.1)}) \\ &= g((p - \alpha \mathbf{1}) \vee q) + g(p \wedge (q + \alpha \mathbf{1})) \quad (\text{by (2.2)}). \end{aligned}$$

## 2.2. Examples

We present several examples of L-convex and  $L^{\sharp}$ -convex functions. See Section 3.3 for more examples.

**Example 2.2** (linear functions). For a real vector  $c \in \mathbb{R}^n$ , the linear function given by  $g(p) = \sum_{i=1}^n c(i)p(i)$  ( $p \in \mathbb{Z}^n$ ) is L-convex as well as  $L^{\sharp}$ -convex. Moreover, given  $a(i, j) \in \mathbb{Z} \cup \{-\infty\}$  ( $i, j \in N, i \leq j$ ) and  $b(i, j) \in \mathbb{Z} \cup \{+\infty\}$  ( $i, j \in N, i \leq j$ ), the function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  given by

$$\begin{aligned} \text{dom } g &= \{p \in \mathbb{Z}^n \mid a(i, i) \leq p(i) \leq b(i, i) \ (i \in N), \\ &\quad a(i, j) \leq p(i) - p(j) \leq b(i, j) \ (i, j \in N, i < j)\}, \\ g(p) &= \sum_{i=1}^n c(i)p(i) \quad (p \in \text{dom } g) \end{aligned}$$

is  $L^{\sharp}$ -convex, provided that  $\text{dom } g$  is nonempty; the function  $g$  is L-convex if  $a(i, i) = -\infty$  ( $i \in N$ ) and  $b(i, i) = +\infty$  ( $i \in N$ ), in addition.  $\square$

**Example 2.3** (quadratic functions). A quadratic function  $g(p) = p^{\top} A p$  ( $p \in \mathbb{Z}^n$ ) given by an  $n \times n$  real symmetric matrix  $A$  is  $L^{\sharp}$ -convex if and only if each off-diagonal component  $a(i, j)$  ( $i, j \in N, i \neq j$ ) is non-positive and the sum of components in each row (or column)  $\sum_{j=1}^n a(i, j)$  ( $i \in N$ ) is non-negative [41].  $\square$

**Example 2.4** (maximum-value functions). Define

$$\begin{aligned} g(p) &= \max\{p(1), p(2), \dots, p(n)\} \quad (p \in \mathbb{Z}^n), \\ g_0(p) &= \max\{0, p(1), p(2), \dots, p(n)\} \quad (p \in \mathbb{Z}^n). \end{aligned}$$

Then,  $g$  is an L-convex function and  $g_0$  is an  $L^{\natural}$ -convex function. We also consider the following functions associated with real numbers  $a(0), a(1), \dots, a(n), b(0), b(1), \dots, b(n) \in \mathbb{R}$ :

$$\begin{aligned}\tilde{g}(p) &= \max_{1 \leq i \leq n} \{p(i) + a(i)\} - \min_{1 \leq i \leq n} \{p(i) + b(i)\} & (p \in \mathbb{Z}^n), \\ \tilde{g}_0(p) &= \max_{0 \leq i \leq n} \{p(i) + a(i)\} - \min_{0 \leq i \leq n} \{p(i) + b(i)\} & (p \in \mathbb{Z}^n),\end{aligned}$$

where  $p(0) = 0$ . Function  $\tilde{g}$  is L-convex and  $\tilde{g}_0$  is  $L^{\natural}$ -convex [37].  $\square$

### 2.3. Properties of minimizers

Minimizers of L-convex and  $L^{\natural}$ -convex functions enjoy various properties. Since an  $L^{\natural}$ -convex function satisfies submodularity (see (2.4)), the set of minimizers forms a sublattice of  $\mathbb{Z}^n$ . We denote by  $\arg \min g$  the set of minimizers of a function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ , i.e.,

$$\arg \min g = \{p \in \mathbb{Z}^n \mid g(p) \leq g(q) \ (\forall q \in \mathbb{Z}^n)\}.$$

**Theorem 2.5.** *Let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^{\natural}$ -convex function.*

- (i) *For  $p, q \in \arg \min g$ , we have  $p \vee q, p \wedge q \in \arg \min g$ .*
- (ii) *Minimal and maximal minimizers of  $g$  are uniquely determined if they exist. In particular, if  $\arg \min g$  is bounded, then minimal and maximal minimizers of  $g$  are uniquely determined.*  $\square$

It should be noted that there may exist many minimal and maximal minimizers for a general function.

The global minimality of an  $L^{\natural}$ -convex function can be characterized by the local minimality with respect to a certain neighborhood. For a function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $p \in \text{dom } g$ , and  $d \in \mathbb{Z}^n$ , we denote by  $g'(p; d)$  the *slope* of  $g$  at  $p$  in the direction  $d$ , i.e.,  $g'(p; d) = g(p + d) - g(p)$ .

**Theorem 2.6** ([31–33]). *Let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^{\natural}$ -convex function and  $p \in \text{dom } g$ . Then, we have  $p \in \arg \min g$  if and only if  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n \cup \{0, -1\}^n$ .*  $\square$

For  $p \in \text{dom } g$ , a vector  $d = d_* \in \{0, +1\}^n \cup \{0, -1\}^n$  minimizing  $g'(p; d)$  is called a *steepest descent direction* of  $g$  at  $p$ . Since  $g'(p; \mathbf{0}) = 0$ , every steepest descent direction  $d_*$  satisfies  $g'(p; d_*) \leq g'(p; \mathbf{0}) = 0$ . Therefore, we have the following equivalence:

$$\begin{aligned}g'(p; d) &\geq 0 \quad (\forall d \in \{0, +1\}^n \cup \{0, -1\}^n) \\ \iff g'(p; d_*) &= 0 \quad (\forall d_* : \text{steepest descent direction at } p) \\ \iff d_* &= \mathbf{0} \text{ is a steepest descent direction at } p.\end{aligned}$$

Due to  $L^{\natural}$ -convexity, minimal and maximal steepest descent directions are uniquely determined.

If we know in advance that a vector  $p \in \text{dom } g$  is a lower bound (or an upper bound) of some minimizer, then we can characterize the global minimality of  $p$  by using the local minimality with respect to a smaller neighborhood.

**Theorem 2.7** (cf. [44]). *Let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^{\natural}$ -convex function, and  $p \in \text{dom } g$  be a vector that is a lower bound of some minimizer  $p_*$  of  $g$  (i.e.,  $p \leq p_*$  holds).*

- (i) *There exists a steepest descent direction  $d$  of  $g$  at  $p$  such that  $d \geq 0$  (i.e.,  $d \in \{0, +1\}^n$ ).*
- (ii)  *$p \in \arg \min g$  holds if and only if  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ .*  $\square$

**Theorem 2.8** (cf. [44]). *Let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^{\natural}$ -convex function, and  $p \in \text{dom } g$  be a vector that is an upper bound of some minimizer  $p_*$  of  $g$  (i.e.,  $p \geq p_*$  holds).*

- (i) *There exists a steepest descent direction  $d$  of  $g$  at  $p$  such that  $d \leq 0$  (i.e.,  $d \in \{0, -1\}^n$ ).*
- (ii)  *$p \in \arg \min g$  holds if and only if  $g'(p; d) \geq 0$  holds for every  $d \in \{0, -1\}^n$ .*  $\square$

For an L-convex function  $g$  that has a minimizer  $p_*$ , the real number  $r$  in the condition (2.2) must be equal to zero, from which follows that  $p_* + \alpha \mathbf{1}$  is also a minimizer of  $g$  for every  $\alpha \in \mathbb{Z}$ . Therefore, every  $p \in \text{dom } g$  is a lower bound of some minimizer as well as an upper bound of another minimizer, which, together with Theorems 2.7 and 2.8, implies the following property.

**Theorem 2.9.** *For an L-convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  and a vector  $p \in \text{dom } g$ , the following three conditions are equivalent:*

- (a)  $p \in \arg \min g$ , (b)  $g'(p; d) \geq 0$  ( $\forall d \in \{0, +1\}^n$ ), (c)  $g'(p; d) \geq 0$  ( $\forall d \in \{0, -1\}^n$ ).  $\square$

### 3. Minimization Algorithms

Minimization algorithms for L-convex and  $L^{\natural}$ -convex functions are presented in this section. Since the minimization of an L-convex function is a special case of the minimization of an  $L^{\natural}$ -convex function, we mainly deal with the latter. Throughout this section, we assume that  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is an  $L^{\natural}$ -convex function with bounded  $\text{dom } g$ .

#### 3.1. Fundamental algorithms

The first algorithm finds a minimizer by iteratively moving a solution along steepest descent directions.

**Algorithm 1** (GREEDY).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  arbitrarily, and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n \cup \{0, -1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, +1\}^n \cup \{0, -1\}^n$  minimizing  $g'(p; d)$ . Set  $p := p + d_*$  and go to Step 1.  $\square$

By Theorem 2.6, the algorithm GREEDY outputs a minimizer of an  $L^{\natural}$ -convex function when it terminates. Since the function value  $g(p)$  decreases strictly in each iteration and  $\text{dom } g$  is bounded, the algorithm GREEDY terminates in a finite number of iterations. Note that the computation of a steepest descent direction in each iteration can be done in polynomial time (in  $n$ ) by reduction to the minimization of submodular set functions  $\rho_p^+, \rho_p^-$  given by

$$\rho_p^+(X) = g'(p; +e_X), \quad \rho_p^-(X) = g'(p; -e_X) \quad (X \subseteq N),$$

where  $e_X \in \{0, +1\}^n$  denotes the characteristic vector of  $X \subseteq N$ ; see [25, 49] for the strongly-polynomial time algorithms for submodular set function minimization.

If a lower bound of a minimizer of  $g$  is available as the initial solution, then an increasing-type steepest descent algorithm can be applied by Theorem 2.7. Here, an ‘‘increasing-type’’ algorithm means that in the algorithm each component of vector  $p$  is monotonically increasing (non-decreasing).

**Algorithm 2** (GREEDYUP).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is a lower bound of some minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, +1\}^n$  minimizing  $g'(p; d)$ . Set  $p := p + d_*$  and go to Step 1.  $\square$

Similarly, if an upper bound of some minimizer of  $g$  is available as the initial solution, then a decreasing-type steepest descent algorithm can be applied by Theorem 2.8.

**Algorithm 3** (GREEDYDOWN).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is an upper bound of some minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, -1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, -1\}^n$  minimizing  $g'(p; d)$ . Set  $p := p + d_*$  and go to Step 1.  $\square$




Figure 2: The behavior of the steepest descent algorithms. Each number associated with each integer lattice point denotes the function value of  $g$  at the point. The shaded region shows the set of minimizers of  $g$ .

**Remark 3.1.** The algorithm GREEDYUP is originally proposed as a minimization algorithm of L-convex functions (see, e.g., [33, Section 10.3.1]). If an L-convex function  $g$  has a minimizer, then for every initial vector  $p_0 \in \text{dom } g$  there exists some minimizer  $p_*$  of  $g$  such that  $p_* \geq p_0$  (see the discussion just before Theorem 2.9). Hence, a minimizer of L-convex function  $g$  can be obtained by GREEDYUP with an arbitrarily chosen initial vector  $p_0$ . Similarly, GREEDYDOWN can also be applied to an L-convex function.  $\square$

**Remark 3.2.** The behavior of vector  $p$  in the algorithm GREEDY applied to an  $L^\natural$ -convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  coincides with the behavior of the original variable vector  $p$  in GREEDYUP applied to L-convex function  $\tilde{g} : \mathbb{Z}^{n+1} \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n + 1$  variables given by (2.3).  $\square$

### 3.2. A numerical example

To illustrate the behavior of the steepest descent algorithms explained above, we consider a numerical example of  $L^\natural$ -convex function  $g : \mathbb{Z}^2 \rightarrow \mathbb{R} \cup \{+\infty\}$  given as follows (see Figure 2):

$$\begin{aligned} \text{dom } g &= \{(p(1), p(2)) \in \mathbb{Z}^2 \mid 0 \leq p(i) \leq 4 \ (i = 1, 2)\}, \\ g(p(1), p(2)) &= \max(0, -p(1) + 2, -p(2) + 1, p(1) - 3, \\ &\quad -p(1) + p(2) - 1, 2p(1) - p(2) - 5) \quad ((p(1), p(2)) \in \text{dom } g). \end{aligned}$$

$L^\natural$ -convexity of this function can be checked by using the translation submodularity (2.4) or discrete-midpoint convexity (2.5) (see Theorem 2.1).

Suppose that the algorithm GREEDY is applied to the function  $g$  with the initial solution  $p_0 = (1, 4)$ . Then, the trajectory of vector  $p$  is given by one of the three dotted arrows in Figure 2 starting from  $(1, 4)$ . In particular, the minimizer found by the algorithm is either of  $(3, 4)$  and  $(2, 3)$ .

We then suppose that the algorithm GREEDY or GREEDYUP is applied to  $g$  with another initial solution  $p_0 = (0, 0)$ . Then, the trajectory of vector  $p$  is given by one of the three dashed arrows in Figure 2 starting from  $(0, 0)$ . In particular, the minimizer found by the algorithm is either of  $(2, 1)$  and  $(2, 2)$ .

### 3.3. Steepest descent algorithms in other research fields

We show that various optimization algorithms proposed independently in other research fields can be seen as steepest descent algorithms in Section 3.1 applied to special  $L^\natural$ -convex (or L-convex) functions.

### 3.3.1. Hassin's algorithm for the minimum cost flow problem

Given a directed graph  $G = (V, E)$ , non-negative edge capacity  $c(e) \in \mathbb{R}_+$  ( $e \in E$ ), and edge cost  $\gamma(e) \in \mathbb{R}$  ( $e \in E$ ), the minimum cost flow problem (minimum cost circulation problem) is formulated as follows:

$$\begin{aligned} \text{Minimize} \quad & \sum_{(u,v) \in E} \gamma(u,v)x(u,v) \\ \text{subject to} \quad & \sum_{v:(u,v) \in E} x(u,v) - \sum_{v:(v,u) \in E} x(v,u) = 0 \quad (u \in V), \\ & 0 \leq x(u,v) \leq c(u,v) \quad ((u,v) \in E). \end{aligned}$$

This is a linear programming problem, and its dual is given as

$$\begin{aligned} \text{Maximize} \quad & g_H(p) \equiv \sum_{(u,v) \in E} c(u,v) \min\{0, p(u) - p(v) + \gamma(u,v)\} \\ \text{subject to} \quad & p(v) \in \mathbb{R} \quad (v \in V). \end{aligned}$$

Hassin's algorithm [21] solves the minimum cost flow problem by finding an optimal solution of the dual problem, instead of solving the primal problem directly.

Assume that edge cost  $\gamma(u,v)$  is integer for each  $(u,v) \in E$ . Then, it can be shown that the dual problem has an integral optimal solution. Hence, we may restrict the values of dual variables  $p(v)$  ( $v \in V$ ) to integers, and the objective function  $g_H$  can be regarded as a function in discrete variables. It is known that this function  $g_H$  is an L-concave function in discrete variables (see [32, 33, 36, 42]). Moreover, Hassin's algorithm coincides with the algorithm GREEDYUP (more precisely, the algorithm GREEDYUPMINIMAL in Section 3.4) applied to L-convex function  $-g_H$  (see [43] for details).

For the minimum cost submodular flow problem, which is a generalization of the minimum cost flow problem, Chung-Tcha [7] proposed an algorithm by generalizing Hassin's algorithm under the integrality assumption for the problem input. This algorithm can also be seen as a (variant of) the algorithm GREEDYUP applied to an L-convex function arising from the dual of the minimum cost submodular flow problem (see [43] for details).

### 3.3.2. Discrete optimization approach in computer vision

Given an undirected graph  $G = (V, E)$  and univariate convex functions  $\varphi_v : \mathbb{Z} \rightarrow \mathbb{R} \cup \{+\infty\}$  ( $v \in V$ ),  $\psi_{uv} : \mathbb{Z} \rightarrow \mathbb{R} \cup \{+\infty\}$  ( $(u,v) \in E$ ), consider the following optimization problem:

$$\begin{aligned} \text{(P):} \quad \text{Minimize} \quad & g_{CV}(p) \equiv \sum_{v \in V} \varphi_v(p(v)) + \sum_{(u,v) \in E} \psi_{uv}(p(v) - p(u)) \\ \text{subject to} \quad & p(v) \in \mathbb{Z} \quad (v \in V). \end{aligned}$$

Here, we say that a univariate function  $\varphi : \mathbb{Z} \rightarrow \mathbb{R} \cup \{+\infty\}$  is convex if  $2\varphi(\alpha) \leq \varphi(\alpha - 1) + \varphi(\alpha + 1)$  holds for every  $\alpha \in \mathbb{Z}$  with  $\varphi(\alpha) < +\infty$ . The objective function  $g_{CV}$  of the problem (P) is an  $L^{\natural}$ -convex function. Moreover, if the first term  $\sum_{v \in V} \varphi_v(p(v))$  is removed from the function  $g_{CV}$ , then the resulting function is an L-convex function (see [32, 33, 36, 42]).

The problem (P) is used to process various tasks in computer vision such as panoramic image stitching [54], image restoration [6], minimization of total variation [9], phase unwrapping in SAR images [4], etc. In these problems, the vertex set  $V$  of the graph  $G$  corresponds to the set of pixels in a given image, and edges of the graph represent the neighborhood relation of pixels. Variable  $p(v)$  represents the label of the pixel  $v$  such as disparity, intensity, etc. The function  $\varphi_v$  for each  $v \in V$  is called the data term and used to evaluate the

label of the pixel  $v$ . The function  $\psi_{uv}$  for  $(u, v) \in E$  is called the smoothness term and used to evaluate the difference of the labels for a pair of adjacent pixels  $u$  and  $v$ . The objective function  $g_{CV}$  of the problem (P) is called an energy function in the context of Markov random field [18], and its minimizer corresponds to a set of labels maximizing the a-posteriori probability.

In computer vision, many algorithms for the problem (P) have been proposed (see, e.g., [4, 27]). In particular, Kolmogorov [27] proposed a general framework of minimization algorithms that includes the algorithms GREEDY, GREEDYUP, and GREEDYDOWN as special cases. An algorithm by Bioucas-Dias–Valadão [4] coincides with the algorithm GREEDYUP applied to the problem (P) without the first term  $\sum_{v \in V} \varphi_v(p(v))$ .

### 3.3.3. Iterative auction in auction theory

Consider an auction with multiple indivisible items. In such an auction, the auctioneer needs to find “appropriate” prices of items as well as an “appropriate” allocation of items to bidders. An ascending auction due to Ausubel [3], which is an algorithm for computing such “appropriate” prices by using the Lyapunov function, can be regarded as an application of the algorithm GREEDYUP to a special  $L^{\natural}$ -convex function. We discuss this topic in more details in Section 6.

### 3.4. Computation of minimal and maximal minimizers

For an  $L^{\natural}$ -convex function with bounded effective domain, minimal and maximal minimizers are uniquely determined by Theorem 2.5. We show that unique minimal and maximal minimizers can be computed by using variants of the algorithms in Section 3.1.

We first consider the minimal minimizer. The minimal minimizer of function  $g$  is the same as the unique minimizer of the function

$$g_{\varepsilon}(p) = g(p) + \varepsilon \sum_{i=1}^n p(i),$$

where  $\varepsilon$  is a sufficiently small positive number. Since the function  $g_{\varepsilon}$  is also an  $L^{\natural}$ -convex function, its minimizer can be computed by the algorithm GREEDYUP, although we need to select the value of  $\varepsilon$  appropriately so that a minimizer of  $g_{\varepsilon}$  is also a minimizer of  $g$ . In fact, we do not need to compute such  $\varepsilon$  explicitly. The behavior of the algorithm GREEDYUP applied to the function  $g_{\varepsilon}$  with a sufficiently small  $\varepsilon > 0$  coincides with the behavior of GREEDYUPMINIMAL to be explained below applied to the original function  $g$ . The difference between GREEDYUPMINIMAL and GREEDYUP is in the choices of the initial solution and a steepest descent direction in each iteration.

**Algorithm 4** (GREEDYUPMINIMAL).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is a lower bound of the minimal minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Find a (unique) minimal minimizer  $d = d_* \in \{0, +1\}^n$  of  $g'(p; d)$ .

Set  $p := p + d_*$  and go to Step 1. □

Note that a minimal minimizer  $d = d_* \in \{0, +1\}^n$  of  $g'(p; d)$  found in Step 2 is uniquely determined due to the submodularity of  $g$ .

**Proposition 3.3.** *For an  $L^{\natural}$ -convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  with bounded  $\text{dom } g$ , the algorithm GREEDYUPMINIMAL outputs the unique minimal minimizer.* □

Based on a similar idea, the algorithm GREEDY can also be used to find the minimal minimizer by modifying the condition of the termination and the choice of a steepest descent direction.

**Algorithm 5** (GREEDYMINIMAL).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  arbitrarily and set  $p := p_0$ .

Step 1: Find a (unique) minimal minimizer  $d = d_* \in \{0, +1\}^n \cup \{0, -1\}^n$  of  $g'(p; d)$ .

Step 2: If  $d_* = \mathbf{0}$ , then output  $p$  and stop. Otherwise, set  $p := p + d_*$  and go to Step 1.  $\square$

**Proposition 3.4.** For an  $L^{\natural}$ -convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  with bounded  $\text{dom } g$ , the algorithm GREEDYMINIMAL outputs the unique minimal minimizer.  $\square$

By modifying the algorithm GREEDYDOWN in a similar way, we can obtain the algorithm GREEDYDOWNMINIMAL that finds the minimal minimizer.

**Remark 3.5.** Iteration of the algorithm GREEDYMINIMAL does not necessarily stop even if the current vector  $p$  is a minimizer of  $g$ . For example, let us consider the  $L^{\natural}$ -convex function  $g$  in Section 3.2. If we apply GREEDYMINIMAL with the initial solution  $p_0 = (1, 4)$ , then the trajectory of  $p$  is given as  $(1, 4) \rightarrow (1, 3) \rightarrow (2, 3) \rightarrow (2, 2) \rightarrow (2, 1)$ , which reaches the minimal minimizer  $(2, 1)$  after passing through the minimizers  $(2, 3)$  and  $(2, 2)$ .  $\square$

We then consider the computation of the maximal minimizer. For a general function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ , a vector  $p$  is a maximal minimizer of  $g$  if and only if  $-p$  is a minimal minimizer of the function  $h : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  defined by  $h(q) = g(-q)$  ( $q \in \mathbb{Z}^n$ ). Moreover, function  $g$  is  $L^{\natural}$ -convex if and only if function  $h$  is  $L^{\natural}$ -convex. Hence, an algorithm for computing the maximal minimizer of an  $L^{\natural}$ -convex function can be easily obtained by modification of an algorithm for the minimal minimizer. For example, we can obtain the following algorithm for the unique maximal minimizer of  $L^{\natural}$ -convex function  $g$  by rewriting the algorithm GREEDYUPMINIMAL applied to  $L^{\natural}$ -convex function  $h$  in terms of the function  $g$ ; the resulting algorithm can be seen as a variant of the algorithm GREEDYDOWN.

**Algorithm 6** (GREEDYDOWNMAXIMAL).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is an upper bound of the maximal minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, -1\}^n$ , then output  $p$  and stop.

Step 2: Find a (unique) maximal minimizer  $d = d_* \in \{0, -1\}^n$  of  $g'(p; d)$ .

Set  $p := p + d_*$  and go to Step 1.  $\square$

**Proposition 3.6.** For an  $L^{\natural}$ -convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  with bounded  $\text{dom } g$ , the algorithm GREEDYDOWNMAXIMAL outputs the unique maximal minimizer.  $\square$

By using similar ideas, we can obtain algorithms GREEDYMAXIMAL and GREEDYUPMAXIMAL for the maximal minimizer as variants of GREEDY and GREEDYUP.

**3.5. Use of long step length**

In all the algorithms explained in Sections 3.1 and 3.4, the current vector  $p$  repeats moving along a certain direction by unit step length. We consider a modification of the algorithms where the vector  $p$  moves along the same steepest descent direction  $d_*$  as far as the value of the slope  $g'(p; d_*)$  in the steepest descent direction does not change.

For example, if this modification is applied to the algorithm GREEDYUP, the following algorithm is obtained. For a vector  $p \in \text{dom } g$  and a direction  $d \in \{0, +1\}^n \cup \{0, -1\}^n$ , we define the value  $\bar{c}(p; d) \in \mathbb{Z} \cup \{+\infty\}$  by

$$\bar{c}(p; d) = \sup\{\lambda \in \mathbb{Z}_+ \mid g(p + \lambda d) - g(p) = \lambda g'(p; d)\}.$$

**Algorithm 7** (GREEDYUP-LS).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is a lower bound of some minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, +1\}^n$  minimizing  $g'(p; d)$ . Set  $\lambda := \bar{c}(p; d_*)$ ,  $p := p + \lambda d_*$ , and go to Step 1.  $\square$

As shown in Section 4.2, this algorithm can be seen as a special implementation of GREEDYUP. In particular, the output of GREEDYUP-LS is a minimizer of  $g$ .

Computation of a steepest descent direction is essentially equivalent to the minimization of a submodular set function (see Section 3.1), for which we require quite large running time to obtain a minimizer (see, e.g., [25, 49]). Therefore, the use of long step length helps to reduce the total running time. Note that the computation of the step length  $\bar{c}(p; d_*)$  can be done by binary search; for the special cases of  $L^{\natural}$ -convex functions shown in Sections 2.2 and 3.3, the step length can be computed more easily.

Long step length can be used to other steepest descent algorithms in a similar way; we omit the details in this paper.

#### 4. Analysis of Number of Iterations in Steepest Descent Algorithms

We analyze the number of iterations required by the algorithms in Section 3. In Section 4.1, the precise bounds for the steepest descent algorithms and its variants are provided in terms of the distance to minimizers. In Section 4.2, we consider an increasing-type steepest descent algorithm using long step length, and obtain an upper bound of the number of iterations by using the slope of steepest descent directions. While other minimization algorithms can be analyzed in a similar way, we omit the details in this paper.

In the following, let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^{\natural}$ -convex function with bounded  $\text{dom } g$ .

##### 4.1. Analysis by using the distance to minimizers

We analyze the number of iterations of algorithms by using the distance from the current vector  $p \in \mathbb{Z}^n$  to minimizers. In the analysis we use two kinds of distance. We define a function  $\hat{\mu} : \mathbb{Z}^n \rightarrow \mathbb{Z}_+ \cup \{+\infty\}$  by

$$\hat{\mu}(p) = \begin{cases} \min\{\|p_* - p\|_{\infty} \mid p_* \in \arg \min g, p_* \geq p\} & (\text{if } \{p_* \in \arg \min g \mid p_* \geq p\} \neq \emptyset), \\ +\infty & (\text{otherwise}). \end{cases} \quad (4.1)$$

That is,  $\hat{\mu}(p)$  is the  $\ell_{\infty}$ -distance between a vector  $p$  and a nearest minimizer  $p_*$  under the condition  $p_* \geq p$ . We also define a function  $\mu : \mathbb{Z}^n \rightarrow \mathbb{Z}_+$  by

$$\mu(p) = \min\{\|p_* - p\|_{\infty}^+ + \|p_* - p\|_{\infty}^- \mid p_* \in \arg \min g\} \quad (p \in \mathbb{Z}^n), \quad (4.2)$$

where

$$\begin{aligned} \|p_* - p\|_{\infty}^+ &= \max_{i \in N} \max(0, p_*(i) - p(i)), \\ \|p_* - p\|_{\infty}^- &= \max_{i \in N} \max(0, -p_*(i) + p(i)). \end{aligned}$$

Function  $\mu(p)$  gives a distance between  $p$  and a nearest minimizer  $p_*$ .

Due to the  $L^{\natural}$ -convexity of  $g$ , the functions  $\mu(p)$  and  $\hat{\mu}(p)$  have the following relationship; the proof is given in Appendix.

**Proposition 4.1** (K. Murota).  $\hat{\mu}(p) = \mu(p)$  holds for a vector  $p \in \mathbb{Z}^n$  such that  $\{q \in \arg \min g \mid q \geq p\} \neq \emptyset$ .  $\square$

**Example 4.2.** Consider the example of an  $L^{\natural}$ -convex function in Section 3.2 (see Figure 2). We have  $\hat{\mu}(p) = \mu(p) = 2$  for  $p = (1, 4)$ . The value  $\hat{\mu}(p) = 2$  is attained by the minimizer  $(3, 4)$ , while  $\mu(p) = 2$  is attained by both of the two minimizers  $(3, 4)$  and  $(2, 3)$ .  $\square$

We first analyze the two increasing-type steepest descent algorithms, GREEDYUP and GREEDYUPMINIMAL. Suppose that a vector  $p \in \text{dom } g$  is a lower bound of some minimizer of function  $g$ . We consider the unique minimal minimizer  $p_*$  of  $g$  under the condition  $p_* \geq p$ , which we denote by  $\hat{p}$ . Recall that in an increasing-type algorithm each component of vector  $p$  increases monotonically. Desirable properties of increasing-type algorithms (for an unknown  $\hat{p}$ ) are as follows:

- (a) for each  $i \in \arg \max_{j \in N} \{\hat{p}(j) - p(j)\}$ , the value of  $p(i)$  is increased if  $\max_{j \in N} \{\hat{p}(j) - p(j)\} > 0$ .
- (b) for each  $i \in N$  with  $\hat{p}(i) - p(i) = 0$ , the value of  $p(i)$  is unchanged.

The next proposition shows that if vector  $p$  is updated by using a steepest descent direction  $d \in \{0, +1\}^n$ , then (a) holds; moreover, (b) also holds if  $d$  is the minimal steepest descent direction, in particular. We denote  $\text{supp}^+(d) = \{i \in N \mid d(i) > 0\}$  for  $d \in \mathbb{Z}^n$ .

**Proposition 4.3** ([44]). *Let  $p \in \text{dom } g$  be a vector that is a lower bound of some minimizer of  $g$ , and  $\hat{p} \in \arg \min g$  be a (unique) minimal minimizer of  $g$  under the condition  $\hat{p} \geq p$ .*

(i) *For every steepest descent direction  $d \in \{0, +1\}^n$  at  $p$ , we have  $\arg \max_{j \in N} \{\hat{p}(j) - p(j)\} \subseteq \text{supp}^+(d)$ . In addition,  $p_* = \hat{p} \vee (p + d)$  is a (unique) minimal minimizer under the condition  $p_* \geq p + d$  and satisfies  $\|p_* - (p + d)\|_\infty = \|\hat{p} - p\|_\infty - 1$ .*

(ii) *For the (unique) minimal steepest descent direction  $d \in \{0, +1\}^n$  at  $p$ , we have*

$$\text{supp}^+(d) \cap \{i \in N \mid \hat{p}(i) = p(i)\} = \emptyset, \quad \hat{p} \geq p + d, \quad \|\hat{p} - (p + d)\|_\infty = \|\hat{p} - p\|_\infty - 1.$$

□

From the proposition above, the precise bounds for the number of updates of  $p$  in GREEDYUP and in GREEDYUPMINIMAL can be obtained. In general, if we update vector  $p$  by increasing each component at most one, then the value  $\hat{\mu}(p)$  decreases by at most one. Hence,  $\hat{\mu}(p_0)$  gives a lower bound for the number of iterations required by any increasing-type algorithm with the initial solution  $p_0$ . In fact, it is shown that the number of updates of  $p$  in GREEDYUP is exactly equal to  $\hat{\mu}(p_0)$ . We define

$$\Phi_g = \max\{\|p - q\|_\infty \mid p, q \in \text{dom } g\},$$

which gives the “size” of the effective domain of  $g$ .

**Theorem 4.4** (cf. [44]).

(i) *If vector  $p$  is updated in an iteration of the algorithm GREEDYUP, then the value  $\hat{\mu}(p)$  decreases by one.*

(ii) *The number of update of vector  $p$  in GREEDYUP is equal to  $\hat{\mu}(p_0)$  and at most  $\Phi_g$ .*

(iii) *The minimizer  $p_* \in \arg \min g$  found by GREEDYUP satisfies  $\|p_* - p_0\|_\infty = \hat{\mu}(p_0)$ .* □

This theorem implies that the trajectory of vector  $p$  generated by GREEDYUP is a shortest path (with respect to  $\ell_\infty$ -distance) from the initial solution  $p_0$  to a nearest minimizer  $p_*$  under the condition  $p_* \geq p_0$ .

**Corollary 4.5** ([44, 46]). *The algorithm GREEDYUPMINIMAL outputs the unique minimal minimizer  $\underline{p}_*$  after vector  $p$  is updated  $\|\underline{p}_* - p_0\|_\infty$  times.* □

**Example 4.6.** We consider the example for the behavior of the algorithm in Section 3.2. If we apply the algorithm GREEDYUP with the initial solution  $p_0 = (0, 0)$ , then the output of the algorithm is either  $p_* = (2, 1)$  or  $p_* = (2, 2)$ , each of which is a nearest minimizer to  $p_0$  (i.e., a minimizer with the minimum value of  $\|p_* - p_0\|_\infty$ ) under the condition  $p_* \geq p_0$ . The number of updates of  $p$  in the algorithm is equal to  $\hat{\mu}(p_0) = 2$ , the distance from  $p_0$  to a nearest minimizer. Three possible trajectories of  $p$  in the algorithm GREEDYUP are drawn by dashed arrows in Figure 2, each of which is a shortest path from the initial

solution  $p_0 = (0, 0)$  to a nearest minimizer. Also, if we apply GREEDYUPMINIMAL, then the trajectory of  $p$  is given as  $(0, 0) \rightarrow (1, 0) \rightarrow (2, 1)$ , and its output  $p_* = (2, 1)$  is the minimal minimizer.  $\square$

A similar result can be obtained for the decreasing-type steepest descent algorithms GREEDYDOWN and GREEDYDOWNMAXIMAL. For a vector  $p \in \text{dom } g$  that is an upper bound of some minimizer of  $L^1$ -convex function  $g$ , we define the value  $\check{\mu}(p) \in \mathbb{Z}_+$  by

$$\check{\mu}(p) = \begin{cases} \min\{\|p_* - p\|_\infty \mid p_* \in \arg \min g, p_* \leq p\} & (\text{if } \{p_* \in \arg \min g \mid p_* \leq p\} \neq \emptyset), \\ +\infty & (\text{otherwise}). \end{cases}$$

**Proposition 4.7** (K. Murota).  $\check{\mu}(p) = \mu(p)$  holds for a vector  $p \in \mathbb{Z}^n$  such that  $\{q \in \arg \min g \mid q \leq p\} \neq \emptyset$ .  $\square$

**Theorem 4.8** (cf. [44]).

(i) If vector  $p$  is updated in an iteration of the algorithm GREEDYDOWN, then the value  $\check{\mu}(p)$  decreases by one.

(ii) The number of update of vector  $p$  in GREEDYDOWN is equal to  $\check{\mu}(p_0)$  and at most  $\Phi_g$ .

(iii) The minimizer  $p_* \in \arg \min g$  found by GREEDYDOWN satisfies  $\|p_* - p_0\|_\infty = \check{\mu}(p_0)$ .  $\square$

**Corollary 4.9** ([44, 46]). The algorithm GREEDYDOWNMAXIMAL outputs the unique maximal minimizer  $\bar{p}_*$  after vector  $p$  is updated  $\|\bar{p}_* - p_0\|_\infty$  times.  $\square$

**Remark 4.10.** We consider the special case where  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is an  $L$ -convex function. If  $g$  has a minimizer, then for every vector  $p_0$ , there exists a minimizer  $p_*$  satisfying  $p_* \geq p_0$  and  $\min_{i \in N} \{p_*(i) - p_0(i)\} = 0$  (see the discussion just before Theorem 2.9). Hence,  $\hat{\mu}(p_0) \leq \Psi_g$  holds with

$$\Psi_g = \max\{\|p - q\|_\infty \mid p, q \in \text{dom } g, \min_{i \in N} \{p(i) - q(i)\} = 0\}.$$

If the value  $\Psi_g$  is finite, then the increasing-type steepest descent algorithms GREEDYUP and GREEDYUPMINIMAL terminate in at most  $\Psi_g$  iterations. Similarly, the decreasing-type steepest descent algorithms GREEDYDOWN and GREEDYDOWNMAXIMAL terminate in at most  $\Psi_g$  iterations.  $\square$

By generalizing the analysis used in Theorem 4.4, we can obtain the exact bound for the number of updates of  $p$  in the algorithms GREEDY and GREEDYMINIMAL.

**Theorem 4.11** ([44]).

(i) If vector  $p$  is updated in an iteration of the algorithm GREEDY, then the value  $\mu(p)$  decreases by one.

(ii) The number of update of vector  $p$  in GREEDY is equal to  $\mu(p_0)$  and at most  $2\Phi_g$ .

(iii) The minimizer  $p_* \in \arg \min g$  found by GREEDY satisfies  $\|p_* - p_0\|_\infty^+ + \|p_* - p_0\|_\infty^- = \mu(p_0)$ .  $\square$

It follows from this theorem that the trajectory of vector  $p$  generated by the algorithm GREEDY is a shortest path (with respect to a certain distance) from the initial solution  $p_0$  to a nearest minimizer  $p_*$ .

**Corollary 4.12** ([44]). The algorithm GREEDYMINIMAL outputs the unique minimal minimizer  $\underline{p}_*$  after vector  $p$  is updated  $\|\underline{p}_* - p_0\|_\infty^+ + \|\underline{p}_* - p_0\|_\infty^-$  times.  $\square$

**Example 4.13.** We reconsider Example 3.2. When we apply the algorithm GREEDY with the initial solution  $p_0 = (1, 4)$ , the output of the algorithm is either of  $p_* = (3, 4)$  and  $p_* = (2, 3)$ , both of which are nearest minimizers to  $p_0$  (with respect to the distance  $\|p_* - p\|_\infty^+ + \|p_* - p\|_\infty^-$ ), and the number of updates of  $p$  is equal to  $\mu(p_0) = 2$ , the distance to a nearest minimizer. Three possible trajectories of  $p$  in the algorithm GREEDY are drawn

by dotted arrows in Figure 2, each of which is a shortest path from the initial solution  $p_0 = (1, 4)$  to a nearest minimizer.  $\square$

**Remark 4.14.** As upper bounds for the number of updates of a vector in the algorithm GREEDY,  $2n\Phi_g$  is shown by Murota [34] and  $2\Phi_g$  by Kolmogorov–Shioura [28]. The result presented in this section (Theorem 4.11) is a refinement of these results by Murota–Shioura [44].  $\square$

**Remark 4.15.** Recently, Hirai [22, 23] introduced the concept of L-convex function on oriented modular graphs, as a generalization of  $L^{\natural}$ -convex function on integer lattice points. In this concept, a function is defined on the set of vertices, and the neighborhood of each vertex is defined by using oriented edges. In this framework, an  $L^{\natural}$ -convex function on integer lattice points can be regarded as an L-convex function on a grid graph with an appropriate edge orientation. Hirai [22, 23] proposed a steepest descent algorithm for minimization of an L-convex function on an oriented modular graph, and showed that the number of iterations is equal to a certain distance between the initial solution and the minimizer output by the algorithm. This result is closely related to Theorems 4.4 and 4.11 presented in this section.  $\square$

#### 4.2. Analysis by the slope of steepest descent direction

We consider a variant of the algorithm GREEDYUPMINIMAL using long step length, which we denote GREEDYUPMINIMAL-LS. In this section, we analyze the number of iterations required by GREEDYUPMINIMAL-LS by using slopes of steepest descent directions. For this, we first show a property concerning the slope of a steepest descent direction in each iteration of GREEDYUPMINIMAL (i.e., the algorithm using the unit step length) and clarify the relationship between GREEDYUPMINIMAL and GREEDYUPMINIMAL-LS.

We show that in GREEDYUPMINIMAL, the absolute value  $|g'(p; d)|$  of the slope of a steepest descent direction  $d$  is non-increasing; moreover, if the unique minimal steepest descent direction is used, then the absolute value  $|g'(p; d)|$  of the slope decreases strictly or the set  $\text{supp}^+(d)$  of non-zero components in a steepest descent direction  $d$  monotonically increases (in the sense of set inclusion). Note that  $g'(p; d) \leq 0$  holds for every  $p \in \text{dom } g$  and every steepest descent direction  $d$  at  $p$ .

**Proposition 4.16** (cf. [16]). *Let  $p \in \text{dom } g$ , and  $d \in \{0, +1\}^n$  be a steepest descent direction at  $p$ , and  $\tilde{d} \in \{0, +1\}^n$  be a steepest descent direction at  $p + d$ .*

(i)  $0 \geq g'(p + d; \tilde{d}) \geq g'(p; d)$  holds.

(ii) Suppose that  $d$  and  $\tilde{d}$  are minimal steepest descent directions at  $p$  and  $p + d$ , respectively. Then, at least one of the following two conditions holds:

- (a)  $g'(p + d; \tilde{d}) > g'(p; d)$ ,      (b)  $g'(p + d; \tilde{d}) = g'(p; d)$  and  $\text{supp}^+(\tilde{d}) \supseteq \text{supp}^+(d)$ .

$\square$

For  $p \in \text{dom } g$ , let  $d \in \{0, +1\}^n$  be a steepest descent direction at  $p$ . Proposition 4.16 (i) implies that  $d$  is also a steepest descent direction at  $p + \lambda d$  for every  $\lambda \in \mathbb{Z}_+$  with  $0 \leq \lambda < \bar{c}(p; d)$ . Hence, the algorithm GREEDYUPMINIMAL-LS using long step lengths can be regarded as a variant of the algorithm GREEDYUPMINIMAL, where the same steepest descent direction as in the previous iteration is used whenever possible. From this observation follows that the trajectory of vector  $p$  of the algorithm GREEDYUPMINIMAL-LS is the same as the trajectory of vector  $p$  of GREEDYUPMINIMAL, and the output of GREEDYUPMINIMAL-LS is a minimizer of function  $g$ .

An upper bound for the number of iterations of GREEDYUPMINIMAL-LS can be obtained by using Proposition 4.16 (ii). When vector  $p$  is updated to  $\tilde{p} = p + \lambda d$  with  $\lambda = \bar{c}(p; d)$ , the slopes  $g'(p; d)$  and  $g'(\tilde{p}; d)$  satisfy  $0 \geq g'(\tilde{p}; d) > g'(p; d)$  by the definition of

$\lambda$  and the discrete mid-point convexity (2.5) of  $g$ . Therefore, Proposition 4.16 (ii) implies that for the unique minimal steepest descent direction  $\tilde{d} \in \{0, +1\}^n$  at  $\tilde{p}$ , at least one of  $g'(\tilde{p}; \tilde{d}) > g'(p; d)$  and  $\text{supp}^+(\tilde{d}) \supsetneq \text{supp}^+(d)$  holds if  $g'(p; d) < 0$ . Since the slope of a steepest descent direction is equal to zero at the termination of the algorithm, the next theorem follows.

**Theorem 4.17.** *For an integer-valued  $L^\natural$ -convex function  $g$ , the number of iterations of the algorithm GREEDYUPMINIMAL-LS is at most  $n \cdot \max\{-g'(p_0; d) \mid d \in \{0, +1\}^n\}$ .  $\square$*

**Remark 4.18.** Consider the example of the minimum cost flow problem in Section 3.3.1. The function  $g_H$  is integer-valued if cost  $c(u, v)$  and capacity  $\gamma(u, v)$  are integer-valued for every edge  $(u, v) \in E$ . Hence, from Theorem 4.17 the following upper bound for the number of iterations of Hassin's algorithm can be obtained:

$$n \cdot \max\{g'_H(p_0; d) \mid d \in \{0, +1\}^n\}.$$

Recall that function  $-g_H$  is an  $L^\natural$ -convex function and its slope is given by  $-g'_H(p; d)$ . This bound coincides with the one shown in Hassin [21, Corollary 3]. Since the slope  $g'_H(p; d)$  at a vector  $p \in \text{dom } g_H$  can be represented as

$$g'_H(p; d) = \sum_{(u,v) \in E'} c(u, v) - \sum_{(u,v) \in E''} c(u, v)$$

for some subsets  $E', E'' \subseteq E$  of edges, we have

$$g'_H(p_0; d) \leq \sum_{(u,v) \in E} |c(u, v)|.$$

From this we obtain an upper bound  $n \cdot \sum_{(u,v) \in E} |c(u, v)|$  for the number of iterations of Hassin's algorithm, which is independent of the choice of the initial solution  $p_0$ .  $\square$

## 5. Minimization of L-convex Function in Continuous Variables

The concepts of L-convexity and  $L^\natural$ -convexity naturally extend to functions in continuous variables. Moreover, the minimization algorithms shown in Section 3 as well as their properties can also be extended to L-convex and  $L^\natural$ -convex functions in continuous variables.

### 5.1. Definition of L-convex function in continuous variables

A convex function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  in continuous variables is called an *L-convex function* [39, 40] if it satisfies the following two properties, where  $\text{dom}_{\mathbb{R}} g = \{p \in \mathbb{R}^n \mid g(p) < +\infty\}$ :

$$\text{[Submodularity]} \quad g(p) + g(q) \geq g(p \vee q) + g(p \wedge q) \quad (\forall p, q \in \text{dom}_{\mathbb{R}} g),$$

**[Linearity in Direction of 1]**

$$\exists r \in \mathbb{R} : g(p + \alpha \mathbf{1}) = g(p) + \alpha r \quad (\forall p \in \mathbb{R}^n, \forall \alpha \in \mathbb{R}).$$

For an L-convex function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ , the function  $\check{g} : \mathbb{R}^{n-1} \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n - 1$  variables given as

$$\check{g}(q(1), \dots, q(n-1)) = g(q(1), \dots, q(n-1), 0)$$

is called an  *$L^\natural$ -convex function* [39, 40]. In other words, a convex function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is said to be  $L^\natural$ -convex if the function  $\tilde{g} : \mathbb{R}^{n+1} \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $n + 1$  variables given by

$$\tilde{g}(q(1), \dots, q(n), q(n+1)) = g(q(1) - q(n+1), \dots, q(n) - q(n+1)) \quad (q \in \mathbb{R}^{n+1})$$

is an L-convex function. As in the case of functions in discrete variables, the function  $\tilde{g}$  defined above satisfies the linearity in the direction of  $\mathbf{1}$ . Hence, L-convexity of function  $\tilde{g}$  is equivalent to submodularity of function  $\tilde{g}$ . This means that  $L^{\natural}$ -convexity of function  $g$  is equivalent to submodularity of function  $\tilde{g}$ .

As in the case of functions in discrete variables,  $L^{\natural}$ -convexity of function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  can be characterized by the translation submodularity:

**[Translation Submodularity]**

$$g(p) + g(q) \geq g((p - \alpha\mathbf{1}) \vee q) + g(p \wedge (q + \alpha\mathbf{1})) \quad (\forall p, q \in \text{dom}_{\mathbb{R}} g, \forall \alpha \in \mathbb{R}_+).$$

**Remark 5.1.** It should be noted that  $L^{\natural}$ -convexity cannot be obtained from the mid-point convexity

$$g(p) + g(q) \geq 2g\left(\frac{p+q}{2}\right) \quad (\forall p, q \in \text{dom}_{\mathbb{R}} g).$$

Under the continuity of a function, the mid-point convexity is equivalent to convexity in the ordinary sense, which is more general than  $L^{\natural}$ -convexity. □

It is clear from their definitions that L-convex and  $L^{\natural}$ -convex functions in continuous variables have close relationship with L-convex and  $L^{\natural}$ -convex functions in discrete variables. If a function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is an L-convex (resp.,  $L^{\natural}$ -convex) function in continuous variables, then its restriction  $g_{\mathbb{Z}} : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  on integer lattice points is an L-convex (resp.,  $L^{\natural}$ -convex) function in discrete variables. On the other hand, in the examples of L-convex (resp.,  $L^{\natural}$ -convex) functions in discrete variables shown in Section 2.2, if we replace the domain  $\mathbb{Z}^n$  with  $\mathbb{R}^n$ , then we can obtain L-convex (resp.,  $L^{\natural}$ -convex) functions in continuous variables. In this way, when we are given L-convex and  $L^{\natural}$ -convex functions in discrete variables, it is often possible to obtain L-convex and  $L^{\natural}$ -convex functions in continuous variables by replacing the domain  $\mathbb{Z}^n$  with  $\mathbb{R}^n$ . Furthermore, it is known that every L-convex (resp.,  $L^{\natural}$ -convex) function in discrete variables can be extended to an L-convex (resp.,  $L^{\natural}$ -convex) functions in continuous variables (see, e.g., [32, 33, 36, 42]).

**5.2. Minimization algorithms and their properties**

Minimization algorithms for  $L^{\natural}$ -convex functions in discrete variables can be naturally extended to polyhedral  $L^{\natural}$ -convex functions. A convex function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is said to be *polyhedral* if its epigraph  $\{(p, \beta) \mid p \in \mathbb{R}^n, \beta \in \mathbb{R}, g(p) \leq \beta\}$  is a polyhedron (i.e., represented as the intersection of a finite number of closed half-spaces). In this section, we assume that  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is a polyhedral  $L^{\natural}$ -convex function.

For  $p \in \text{dom}_{\mathbb{R}} g$  and  $d \in \mathbb{R}^n$ , the *directional derivative*  $g'(p; d)$  of function  $g$  at  $p$  in the direction  $d$  is given as

$$g'(p; d) = \lim_{\lambda \downarrow 0} \frac{g(p + \lambda d) - g(p)}{\lambda}.$$

Since  $g$  is a polyhedral convex function, the limit in the right-hand side exists if we allow to have  $g'(p; d) = +\infty$ .

By its definition, an  $L^{\natural}$ -convex function  $g$  in continuous variables is a convex function in the ordinary sense. Hence, a vector  $p \in \text{dom}_{\mathbb{R}} g$  is a minimizer if and only if  $g'(p; d) \geq 0$  holds for every direction  $d \in \mathbb{R}^n$ . In fact, by  $L^{\natural}$ -convexity we may restrict our attention only to a finite number of directions to check the minimality of a vector.

**Theorem 5.2** ([32, 33, 39]). *Let  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be a polyhedral  $L^{\natural}$ -convex function. Then, a vector  $p \in \text{dom}_{\mathbb{R}} g$  is a minimizer of  $g$  if and only if  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n \cup \{0, -1\}^n$ .* □

This is an extension of Theorem 2.6 for  $L^{\natural}$ -convex functions in discrete variables. By Theorem 5.2, the steepest descent algorithm GREEDY-LS using (long) step length in Section 3 can be naturally extended to a polyhedral  $L^{\natural}$ -convex function  $g$ . For  $p \in \text{dom}_{\mathbb{R}} g$  and  $d \in \mathbb{R}^n$  with  $g'(p; d) < +\infty$ , we define the value  $\bar{c}(p; d) \in \mathbb{R} \cup \{+\infty\}$  by

$$\bar{c}(p; d) = \sup\{\lambda \in \mathbb{R}_+ \mid g(p + \lambda d) - g(p) = \lambda g'(p; d)\}. \quad (5.1)$$

Since  $g$  is a polyhedral convex function, the value  $\bar{c}(p; d)$  is always positive, and  $g(p + \lambda d) - g(p) = \lambda g'(p; d)$  holds with  $\lambda = \bar{c}(p; d)$ .

**Algorithm 8** (GREEDY-LS $[\mathbb{R}]$ ).

Step 0: Select an initial solution  $p_0 \in \text{dom}_{\mathbb{R}} g$  arbitrarily and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n \cup \{0, -1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, +1\}^n \cup \{0, -1\}^n$  minimizing  $g'(p; d)$ . Set  $p := p + \lambda d_*$  with  $\lambda := \bar{c}(p; d_*)$  and go to Step 1.  $\square$

Theorem 4.11 for the case of discrete variables can be extended to this algorithm. In a similar way as in Section 4.1, we define a function  $\mu : \mathbb{R}^n \rightarrow \mathbb{R}_+$  by

$$\mu(p) = \min\{\|p_* - p\|_{\infty}^+ + \|p_* - p\|_{\infty}^- \mid p_* \in \arg \min_{\mathbb{R}} g\},$$

where

$$\arg \min_{\mathbb{R}} g = \{p \in \mathbb{R}^n \mid g(p) \leq g(q) \ (\forall q \in \mathbb{R}^n)\}.$$

**Theorem 5.3** ([16]). *Let  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be a polyhedral  $L^{\natural}$ -convex function such that a minimizer exists. In each iteration of the algorithm GREEDY-LS $[\mathbb{R}]$ , the value  $\mu(p)$  decreases by the step length  $\bar{c}(p; d)$ . In addition, the algorithm outputs a minimizer  $p_*$  of  $g$  satisfying  $\|p_* - p_0\|_{\infty}^+ + \|p_* - p_0\|_{\infty}^- = \mu(p_0)$  when it terminates.*  $\square$

This theorem implies that when the algorithm GREEDY-LS $[\mathbb{R}]$  is applied to a polyhedral  $L^{\natural}$ -convex function, its output is a minimizer  $p_*$  nearest to the initial solution, and the trajectory of vector  $p$  is a shortest path from the initial solution to  $p_*$ . It should be noted that this theorem does not guarantee the termination of the algorithm.

Theorem 2.7 (ii) for the case of discrete variables can also be extended to polyhedral  $L^{\natural}$ -convex functions as follows.

**Theorem 5.4** (cf. [44]). *Let  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be a polyhedral  $L^{\natural}$ -convex function such that a minimizer exists, and  $p \in \text{dom}_{\mathbb{R}} g$  be a vector that is a lower bound of some minimizer of  $g$ . Then,  $p$  is a minimizer of  $g$  if and only if  $g'(p; d) \geq 0$  for every  $d \in \{0, +1\}^n$ .*  $\square$

By this theorem the increasing-type steepest descent algorithm GREEDYUP-LS can also be extended to a polyhedral  $L^{\natural}$ -convex function  $g$ .

**Algorithm 9** (GREEDYUP-LS $[\mathbb{R}]$ ).

Step 0: Select an initial solution  $p_0 \in \text{dom}_{\mathbb{R}} g$  that is a lower bound of some minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Find  $d = d_* \in \{0, +1\}^n$  minimizing  $g'(p; d)$ . Set  $p := p + \lambda d_*$  with  $\lambda := \bar{c}(p; d_*)$  and go to Step 1.  $\square$

Theorem 4.4 for the case of discrete variables can be extended to this algorithm. As in Section 4.1, we define a function  $\hat{\mu} : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$\hat{\mu}(p) = \begin{cases} \min\{\|p_* - p\|_{\infty} \mid p_* \in \arg \min_{\mathbb{R}} g, p_* \geq p\} & (\text{if } \{p_* \in \arg \min_{\mathbb{R}} g \mid p_* \geq p\} \neq \emptyset), \\ +\infty & (\text{otherwise}). \end{cases}$$

**Theorem 5.5** (cf. [16]).

(i) If vector  $p$  is updated in an iteration of the algorithm GREEDYUP-LS[ $\mathbb{R}$ ], then the value  $\hat{\mu}(p)$  decreases by the step length  $\bar{c}(p; d)$ .

(ii) The minimizer  $p_* \in \arg \min g$  found by GREEDYUP-LS[ $\mathbb{R}$ ] satisfies  $\|p_* - p_0\|_\infty = \hat{\mu}(p_0)$ . □

Theorems 5.2 and 5.4 imply that the algorithms GREEDY-LS[ $\mathbb{R}$ ] and GREEDYUP-LS[ $\mathbb{R}$ ] output minimizers when they terminate. These algorithms, however, have no guarantee to terminate in a finite number of iterations; furthermore, there exists a polyhedral  $L^\natural$ -convex function for which the sequence of vector  $p$  generated by each of the algorithm converges to a vector that is not a minimizer. Indeed, it is shown in [16] that for some concrete example of the function  $-g_H$  in Section 3.3.1 neither GREEDYUP-LS[ $\mathbb{R}$ ] terminates nor the sequence of vector  $p$  in the algorithm converges to a minimizer; recall that  $-g_H$  is a polyhedral L-convex function.

On the other hand, we can show that the algorithm terminates in a finite number of iterations if we use a special choice of a steepest descent direction in each iteration. For example, use of minimal steepest descent directions in GREEDYUP-LS[ $\mathbb{R}$ ] guarantees the finite termination.

**Algorithm 10** (GREEDYUPMINIMAL-LS[ $\mathbb{R}$ ]).

Step 0: Select an initial solution  $p_0 \in \text{dom } g$  that is a lower bound of the minimal minimizer of  $g$ , and set  $p := p_0$ .

Step 1: If  $g'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Find a (unique) minimal minimizer  $d = d_* \in \{0, +1\}^n$  of  $g'(p; d)$ . Set  $p := p + \lambda d_*$  with  $\lambda := \bar{c}(p; d_*)$  and go to Step 1. □

For this algorithm a property similar to Proposition 4.16 (ii) for  $L^\natural$ -convex function in discrete variables holds.

**Proposition 5.6** (cf. [16]). *Let  $p \in \text{dom}_{\mathbb{R}} g$ , and  $d \in \{0, +1\}^n$  be the unique minimal minimizer of  $g'(p; d)$ . Also, let  $\tilde{p} = p + \bar{c}(p; d)d$ , and  $\tilde{d} \in \{0, +1\}^n$  be the unique minimal minimizer of  $g'(\tilde{p}; \tilde{d})$ . Then, we have  $g'(\tilde{p}; \tilde{d}) \geq g'(p; d)$ . Moreover, if  $g'(\tilde{p}; \tilde{d}) = g'(p; d) < 0$  then we have  $\text{supp}^+(\tilde{d}) \supsetneq \text{supp}^+(d)$ .* □

For a polyhedral convex function  $g$ , the set of possible values of the directional derivatives  $g'(p; d)$  for  $p \in \text{dom}_{\mathbb{R}} g$  and  $d \in \{0, +1\}^n$  is finite. This fact, together with the proposition above, implies the finite termination of GREEDYUPMINIMAL-LS[ $\mathbb{R}$ ].

**Theorem 5.7** (cf. [16]). *For a polyhedral  $L^\natural$ -convex function  $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  such that a minimizer exists, the algorithm GREEDYUPMINIMAL-LS[ $\mathbb{R}$ ] terminates in a finite number of iterations and outputs a minimizer of  $g$ . Moreover, the algorithm outputs the unique minimal minimizer of  $g$  if the initial solution  $p_0$  is a lower bound of the minimal minimizer.* □

## 6. Application to Auction Theory

In this section, we show that results on L-convex function minimization algorithms explained so far can be applied to iterative auctions in auction theory. Iterative auctions are used to compute equilibrium prices of goods. It is shown that iterative auctions for a certain general model can be seen as minimization algorithms applied to a special  $L^\natural$ -convex function.

### 6.1. Model of auction and fundamental concepts

The auction model and fundamental concepts used in this section are explained below. Let us consider an auction with  $n$  types of indivisible goods, where the set of goods is denoted as  $N = \{1, 2, \dots, n\}$ . We suppose that  $m$  bidders attend the auction, and denote the set

of bidders by  $M = \{1, 2, \dots, m\}$ . Each bidder  $j \in M$  has his/her own valuation function  $f_j : 2^N \rightarrow \mathbb{R}$ , where the value  $f_j(X)$  represents the degree of satisfactory for the set of goods  $X \subseteq N$ . Given a price vector  $p \in \mathbb{R}_+^n$ , each bidder  $j \in M$  wants to buy a set of goods  $X_j$  that maximizes the utility  $f_j(X_j) - p(X_j)$ . For each bidder  $j \in M$ , we denote by  $V_j(p) \in \mathbb{R}$  the maximum utility of the bidder  $j$  and by  $D_j(p) \subseteq 2^N$  a family of sets of goods maximizing utility. That is,

$$V_j(p) = \max\{f_j(X) - \sum_{i \in X} p(i) \mid X \subseteq N\} \quad (p \in \mathbb{R}_+^n), \quad (6.1)$$

$$D_j(p) = \arg \max\{f_j(X) - \sum_{i \in X} p(i) \mid X \subseteq N\} \quad (p \in \mathbb{R}_+^n). \quad (6.2)$$

Function  $V_j$  is called the *indirect utility function* of bidder  $j$ , and the set family  $D_j(p)$  for  $p \in \mathbb{R}_+^n$  is called the *demand set* (or *demand correspondence*) of bidder  $j$ .

The goal of an auction is to find a partition  $\{X_1^*, X_2^*, \dots, X_m^*\}$  (i.e., an allocation of goods to bidders) of goods in  $N$  and a price vector  $p_*$  satisfying the condition  $X_j^* \in D_j(p_*)$  ( $j \in M$ ). That is, we want to find an allocation of goods and a price vector such that each bidder is fully satisfied and all goods are sold out. Such a pair of a partition of goods  $\{X_1^*, X_2^*, \dots, X_m^*\}$  and a price vector  $p_*$  is called a *Walrasian equilibrium* (or *competitive equilibrium*);  $p_*$  is called a *Walrasian equilibrium price vector*.

While Walrasian equilibrium may not exist in general, it exists if each valuation function  $f_j$  satisfies a natural condition called the gross substitutes condition [19, 26]. The *gross substitutes condition* for a valuation function  $f_j$  is described in terms of demand sets as follows:

$$\text{(GS)} \quad \forall p, q \in \mathbb{R}_+^n \text{ with } p \leq q, \forall X \in D_j(p), \exists Y \in D_j(q) \\ \text{such that } Y \supseteq \{i \in X \mid p(i) = q(i)\}.$$

The gross substitutes condition means that when prices of some goods increase, the goods with increased prices are the only goods that may drop from the demand set. The gross substitutes condition is introduced by Kelso–Crawford [26] in the setting of a fairly general two-sided job matching model. Since then, this condition has been widely used in various models such as matching, housing, and labor markets (see, e.g., [3, 5, 8, 19, 20]).

An *iterative auction* is an algorithm (more precisely, a protocol) to compute an equilibrium by repeatedly updating a price vector [5, 8]. An iterative auction is called an *ascending auction* if prices increase monotonically in each iteration. An ascending auction for multiple goods is a generalization of the well-known English auction for a single good (see, e.g., [5, 8]). In addition, an ascending auction is natural from the economic viewpoint, and easy to understand and to be implemented.

For an auction model with gross substitutes valuation functions, an ascending auction is proposed by Gul–Stacchetti [20]. Later, Ausubel [3] featured the concept of Lyapunov function, which is a function in a price vector given by

$$L(p) = \sum_{j=1}^m V_j(p) + \sum_{i=1}^n p(i) \quad (p \in \mathbb{R}_+^n). \quad (6.3)$$

Ausubel [3] showed that the Lyapunov function enjoys various nice properties.

**Theorem 6.1** ([3]). *Suppose that each valuation function  $f_j$  satisfies the gross substitutes condition.*

(i) *A vector  $p \in \mathbb{R}_+^n$  is a minimizer of the Lyapunov function if and only if it is an equilibrium*

price vector.

(ii) Suppose that each  $f_j$  is integer-valued. Then, the Lyapunov function has an integral minimizer. In particular, a minimal and maximal minimizer of the Lyapunov function are integral vectors.  $\square$

Based on these facts, Ausubel [3] proposed ascending and descending auctions that find an equilibrium price vector through the minimization of the Lyapunov function. Note that the ascending auction of Ausubel [3] can be seen as a reformulation of the one by Gul–Stacchetti [20].

## 6.2. Analysis of iterative auctions

### 6.2.1. Connection between auction theory and discrete convex analysis

We discuss the connection between auction theory and discrete convex analysis in this section. In discrete convex analysis, another class of discrete convex functions called  $M^\natural$ -concave functions [38], in addition to  $L^\natural$ -convex functions, plays a primary role. Similar to  $L^\natural$ -convexity,  $M^\natural$ -concavity is defined for functions on integer lattice points. The definition of  $M^\natural$ -concavity, when specialized to a set function  $f : 2^N \rightarrow \mathbb{R}$ , is described as follows:

a function  $f : 2^N \rightarrow \mathbb{R}$  is called an  $M^\natural$ -concave function if for every  $X, Y \subseteq N$  and every  $u \in X \setminus Y$ , at least one of (a) and (b) holds:

(a)  $f(X) + f(Y) \leq f(X - u) + f(Y + u)$ ,

(b)  $\exists v \in Y \setminus X$  such that  $f(X) + f(Y) \leq f(X - u + v) + f(Y + u - v)$ ,

where  $X - u + v = (X \setminus \{u\}) \cup \{v\}$ .

The connection between auction theory and discrete convex analysis is initiated by the following theorem due to Fujishige–Yang [17]:

**Theorem 6.2** ([17]). *A function  $f : 2^N \rightarrow \mathbb{R}$  satisfies the gross substitutes condition if and only if it is an  $M^\natural$ -concave function.*  $\square$

Based on this theorem, known results in discrete convex analysis can be used in various fields of mathematical economics such as auction theory and game theory.

The following conjugacy relation holds between  $M^\natural$ -concavity and  $L^\natural$ -convexity. We say that a polyhedral  $L^\natural$ -convex function  $g$  is *domain-integral* if the set  $\arg \min\{g(p) - p^\top x \mid p \in \mathbb{R}^n\}$  is an empty set or an integral polyhedron for every  $x \in \mathbb{R}^n$ .

**Theorem 6.3** ([31, 33, 39]). *For a set function  $f : 2^N \rightarrow \mathbb{R}$ , define a function  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  by*

$$g(p) = \max\{f(X) - \sum_{i \in X} p(i) \mid X \subseteq N\}.$$

(i)  *$f$  is an  $M^\natural$ -concave function if and only if  $g$  is a polyhedral  $L^\natural$ -convex function.*

(ii) *Suppose that  $f$  is an integer-valued  $M^\natural$ -concave function. Then,  $g$  is a domain-integral polyhedral  $L^\natural$ -convex function.*  $\square$

$L^\natural$ -convexity of the indirect utility function and the Lyapunov function can be obtained from Theorems 6.2 and 6.3 and the fact that  $L^\natural$ -convexity of a polyhedral convex function is closed under addition.

**Theorem 6.4** ([45, 46]). *Suppose that the valuation function  $f_j : 2^N \rightarrow \mathbb{R}$  of each bidder  $j \in M$  satisfies the gross substitutes condition. Then, for each  $j \in M$ , the indirect utility function  $V_j : \mathbb{R}_+^n \rightarrow \mathbb{R}$  and the Lyapunov function  $L : \mathbb{R}_+^n \rightarrow \mathbb{R}$  are polyhedral  $L^\natural$ -convex functions. Moreover, if each  $f_j$  is an integer-valued function, then the Lyapunov function  $L$  is a domain-integral polyhedral  $L^\natural$ -convex function.*  $\square$

Thus, an iterative auction using the Lyapunov function can be seen as a minimization algorithm for a special  $L^\natural$ -convex function.

### 6.2.2. Ascending and descending auctions

Murota–Shioura–Yang [45, 46] analyzed the ascending and descending auctions of Ausubel [3] from the viewpoint of discrete convex analysis, which is explained in this section. In the following sections, we assume that the valuation functions  $f_j$  are integer-valued and satisfy the gross substitutes condition.

The ascending auction, which finds the unique minimal (integral) equilibrium price vector  $\underline{p}^*$ , can be described as follows:

**Algorithm 11** (ASCENDAUCTION).

Step 0: Select an initial price vector  $p_0 \in \mathbb{Z}_+^n$  with  $p_0 \leq \underline{p}^*$ , and set  $p := p_0$ .

Step 1: If  $L'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ , then output  $p$  and stop.

Step 2: Let  $d = d_* \in \{0, +1\}^n$  be a (unique) minimal minimizer of  $L'(p; d)$ .

Set  $p := p + d_*$  and go to Step 1. □

The algorithm ASCENDAUCTION can be interpreted in auction terms as follows:

Step 0: The auctioneer sets the initial price vector  $p := p^\circ$ , where  $p^\circ \in \mathbb{Z}^n$  satisfies  $p^\circ \leq \underline{p}^*$ .

Step 1: The auctioneer asks the bidders to report their demand sets  $D_j(p)$  ( $j \in M$ ),

and by using the information of demand sets, checks if  $L'(p; d) \geq 0$  holds for every  $d \in \{0, +1\}^n$ . If it holds, the auctioneer reports  $p$  as the final price vector and stop.

Step 2: The auctioneer finds a (unique) minimal minimizer  $d = d_* \in \{0, +1\}^n$  of  $L(p + d)$ , sets  $p := p + d_*$ , and returns to Step 1. □

**Remark 6.5.** We discuss an implementation issue of iterative auctions. It is often the case that values of valuation functions  $f_j$  are not explicitly given since they are bidders' private information. In such a case, the value of the Lyapunov function cannot be computed. The ascending auction of Ausubel [3] uses the information of demand sets  $D_j(p)$  for a given price vector  $p \in \mathbb{Z}_+^n$ , by which the slope  $L'(p; d)$  of the Lyapunov function can be computed for any given direction  $d \in \{0, +1\}^n \cup \{0, -1\}^n$  (see [3, Appendix B] or the technical report version of [45]). Therefore, in the ascending auction it is possible to check the local optimality of the Lyapunov function in Step 1 and to compute a steepest descent direction in Step 2, without using the value of the Lyapunov function. □

**Remark 6.6.** The ascending auction of Ausubel [3] coincides with the classical English auction if the number of goods  $n$  is equal to one. In other words, the ascending auction is a natural generalization of the English auction. □

Ausubel [3] showed that the algorithm ASCENDAUCTION terminates in a finite number of iterations by outputting the unique minimal equilibrium price vector  $\underline{p}_*$ . The exact bound for the number of iterations can be obtained from the results in Section 4.1.

By Theorem 6.4, the Lyapunov function is a polyhedral  $L^{\natural}$ -convex function. Since the minimal equilibrium price vector is an integral vector, we can restrict the Lyapunov function on the integer lattice points; the resulting function can be seen as an  $L^{\natural}$ -convex function in discrete variables. Therefore, we see that the algorithm ASCENDAUCTION coincides with the algorithm GREEDYUPMINIMAL applied to the Lyapunov function. The exact bound for the number of iterations required by ASCENDAUCTION can be obtained from this observation and Corollary 4.5.

**Theorem 6.7** ([45, 46]). *The algorithm ASCENDAUCTION updates a price vector exactly  $\|\underline{p}^* - p^\circ\|_\infty$  times and then terminates by outputting the unique minimal equilibrium price vector  $\underline{p}_*$ .* □

Any ascending auction algorithm requires at least  $\|\underline{p}^* - p^\circ\|_\infty$  iterations if the algorithm increases the price of each good by at most one. This means that Ausubel's ascending

auction finds the minimal equilibrium price vector using the smallest number of iterations, i.e., it is the “fastest” auction algorithm among all ascending auction algorithms.

**Remark 6.8.** In the case where the valuation functions  $f_j$  are real-valued, we can compute an equilibrium price vector *exactly* by applying the algorithm GREEDYUPMINIMAL-LS[ $\mathbb{R}$ ] in Section 5 to the Lyapunov function, provided that the step length  $\bar{c}(p; d_*)$  can be computed *exactly* in each iteration. The exact computation of the step length, however, is quite difficult (or impossible) in practice. Instead, we can compute an *approximate* equilibrium price vector if we can compute the step length *approximately* in each iteration; see the technical report version of [45] for details.  $\square$

The descending auction of Ausubel [3] described below is quite similar to the ascending auction, and iteratively decreases prices and finds the unique maximal equilibrium price vector  $\bar{p}^*$ .

**Algorithm 12** (DESCENDAUCTION).

Step 0: Select an initial price vector  $p_0 \in \mathbb{Z}_+^n$  with  $p_0 \geq \bar{p}^*$ , and set  $p := p_0$ .

Step 1: If  $L'(p; d) \geq 0$  holds for every  $d \in \{0, -1\}^n$ , then output  $p$  and stop.

Step 2: Let  $d = d_* \in \{0, -1\}^n$  be a (unique) maximal minimizer of  $L'(p; d)$ .

Set  $p := p + d_*$  and go to Step 1.  $\square$

**Remark 6.9.** The descending auction of Ausubel [3] coincides with the classical Dutch auction if the number of goods  $n$  is equal to one. In other words, the descending auction is a natural generalization of Dutch auction.  $\square$

The algorithm DESCENDAUCTION coincides with the algorithm GREEDYDOWNMAXIMAL applied to the Lyapunov function. Hence, the exact bound for the number of iterations of DESCENDAUCTION can be obtained in a similar way as ASCENDAUCTION.

**Theorem 6.10** ([45, 46]). *The algorithm DESCENDAUCTION updates a price vector exactly  $\|p^\circ - \bar{p}^*\|_\infty$  times and then terminates by outputting the unique maximal equilibrium price vector  $\bar{p}_*$ .*  $\square$

**Remark 6.11.** A variant of auction algorithm in Gul–Stacchetti [20] uses line search to compute an equilibrium price vector (see [20, p. 77]). It can be shown that the behavior of the auction algorithm coincides with that of the increasing-type steepest descent algorithm GREEDYUPMINIMAL-LS using long step length applied to the Lyapunov function. Hence, we can apply Theorem 4.17 to obtain the number of iterations of the auction algorithm of Gul–Stacchetti [20].

Recall that the Lyapunov function is an integer-valued function under the assumption that each  $f_j$  is an integer-valued function. By the definition of the Lyapunov function (6.3), the slope of the Lyapunov function can be represented as

$$L'(p; d) = - \sum_{j=1}^m d(X_j) + d(N) \quad (d \in \{0, +1\}^n)$$

with some subsets  $X_1, X_2, \dots, X_m$  of  $N$ . Therefore we have  $L'(p; d) \geq -(m - 1)n$ . Hence, the auction algorithm of Gul–Stacchetti [20] terminates at most  $(m - 1)n^2$  iterations; this bound is also shown in Saito [48].  $\square$

### 6.2.3. Greedy auction and two-phase auction

We explain the new iterative auction algorithms proposed by Murota–Shioura–Yang [45, 46]. An important advantage of ascending and descending auctions is that the change of a price vector is monotone. On the other hand, the initial price vector of ascending (resp., descending) auctions should be a lower (resp., upper) bound of an equilibrium price vector,

which should be sufficiently small (resp., large) since an equilibrium price vector is not known in advance. This is a disadvantage of ascending and descending auctions since it makes the number of iterations of the algorithm fairly large.

To eliminate the disadvantage, Murota–Shioura–Yang [45, 46] proposed two kinds of auction algorithms, in both of which any price vector can be used as an initial vector. The first algorithm, called the greedy auction, is an iterative auction obtained by applying GREEDYMINIMAL in Section 3.4 to the Lyapunov function.

**Algorithm 13** (GREEDYAUCTION).

Step 0: Select an initial price vector  $p_0 \in \mathbb{Z}_+^n$  arbitrarily, and set  $p := p_0$ .

Step 1: Let  $d = d_* \in \{0, +1\}^n \cup \{0, -1\}^n$  be a (unique) minimal minimizer of  $L'(p; d)$ .

Step 2: If  $d_* = \mathbf{0}$  then output  $p$  and stop. Otherwise, set  $p := p + d_*$  and go to Step 1.  $\square$

**Remark 6.12.** The greedy auction is a slight modification of the one in Murota–Shioura–Yang [45] so that the minimal equilibrium price vector can be found.  $\square$

The greedy auction has an advantage that any price vector can be used initially. Therefore, if a good estimate of an equilibrium price vector is available from the output of auctions in the past, we can use it as the initial price vector, resulting in the reduction of the number of iterations. Corollary 4.12 for the algorithm GREEDYMINIMAL implies that the number of iterations of GREEDYAUCTION is equal to a distance between the initial price vector  $p_0$  and the minimal equilibrium price vector  $\underline{p}_*$ .

**Theorem 6.13** ([45, 46]). *The algorithm GREEDYAUCTION updates a price vector exactly  $\|\underline{p}_* - p_0\|_\infty^+ + \|\underline{p}_* - p_0\|_\infty^-$  times and then terminates by outputting the unique minimal equilibrium price vector  $\underline{p}_*$ .*  $\square$

The algorithm GREEDYAUCTION, however, has a disadvantage that the change of a price vector is not monotone. On the other hand, the two-phase auction described below has an additional advantage that changes of prices are almost monotone. As is clear from its name, the two-phase auction consists of two phases, the ascending phase and the descending phase.

**Algorithm 14** (TWO PHASE AUCTION).

Step 0: Select an initial price vector  $p_0 \in \mathbb{Z}_+^n$  arbitrarily, set  $p := p_0$ , and go to Ascending Phase.

[Ascending Phase]

Step A1: Let  $d = d_* \in \{0, +1\}^n$  be a (unique) minimal minimizer of  $L'(p; d)$ .

Step A2: If  $d_* = \mathbf{0}$  then go to Descending Phase.

Otherwise, set  $p := p + d_*$  and go to Step A1.

[Descending Phase]

Step D1: Let  $d = d_* \in \{0, -1\}^n$  be a (unique) minimal minimizer of  $L'(p; d)$ .

Step D2: If  $d_* = \mathbf{0}$  then output  $p$  and stop.

Otherwise, set  $p := p + d_*$  and go to Step D1.  $\square$

While the two-phase auction has advantages as mentioned above, the number of iterations can be larger than that of the greedy auction. We can show that the number of iterations of the two-phase auction is at most three times of that of greedy auction.

**Theorem 6.14** ([45, 46]). *The algorithm TWO PHASE AUCTION terminates with outputting the unique minimal equilibrium price vector  $\underline{p}_*$ . The number of updates of a price vector is at most  $\|\underline{p}_* - p_0\|_\infty^+ + \|\underline{p}_* - p_0\|_\infty^-$  in the ascending phase and at most  $2(\|\underline{p}_* - p_0\|_\infty^+ + \|\underline{p}_* - p_0\|_\infty^-)$  in the descending phase.*  $\square$

**Remark 6.15.** The *unit-demand auction model* is a special case of the auction model considered in this section, where each bidder is interested in getting at most one item.

Iterative auctions for this model are discussed extensively in the literature; e.g., Vickrey–English auction [10], Vickrey–Dutch auction [29], and Vickrey–English–Dutch auction [2]. It is shown in [46, Section 5] that these iterative auctions coincide with the ascending auction, the descending auction, and the two-phase auction, respectively, explained in this section applied to unit-demand valuation functions. Note that a unit-demand valuation function is a special case of a gross substitutes valuation function.  $\square$

**Remark 6.16.** Throughout Section 6 we have focused on computing an equilibrium price vector. We here briefly explain how to compute an allocation in an equilibrium, provided that an equilibrium price vector is available.

Given an equilibrium price vector  $p^*$ , we want to find a partition  $\{X_1^*, X_2^*, \dots, X_m^*\}$  of  $N$  satisfying the condition  $X_j^* \in D_j(p^*)$  ( $j \in M$ ). It is known (see, e.g., [33]) that for a valuation function  $f_j$  satisfying the gross substitutes condition, the demand set  $D_j(p^*)$  has a nice combinatorial structure called a *generalized matroid* [52], a generalization of the concept of matroid. Based on this fact, we can reduce the problem of computing an equilibrium allocation to a variant of the matroid union problem, which can be solved by a matroid intersection algorithm (see, e.g., [50]).  $\square$

### Acknowledgement

The author thanks Kazuo Murota for giving valuable comments on the manuscript and for communicating Propositions 4.1 and 4.7. The author also thanks the anonymous referees for careful reading of the manuscript and useful suggestion. This work is supported by JSPS/MEXT KAKENHI Grand Numbers 26280004, 15K00030, 15H00848.

### A. Appendix: Proof of Proposition 4.1

Let  $p_* \in \arg \min g$  be the minimal minimizer under the condition  $p_* \geq p$ ; such  $p_*$  is uniquely determined and satisfies  $\|p_* - p\|_\infty = \hat{\mu}(p)$ . By the definitions of  $\hat{\mu}(p)$  and  $\mu(p)$ , we have

$$\hat{\mu}(p) = \|p_* - p\|_\infty = \|p_* - p\|_\infty^+ = \|p_* - p\|_\infty^+ + \|p_* - p\|_\infty^- \geq \mu(p). \quad (\text{A.1})$$

To show that the inequality (A.1) holds with equality, we assume, to the contrary, that  $\|p_* - p\|_\infty^+ + \|p_* - p\|_\infty^- > \mu(p)$  and derive a contradiction.

Let  $q_* \in \arg \min g$  be a minimizer with  $\|q_* - p\|_\infty^+ + \|q_* - p\|_\infty^- = \mu(p)$ ; we assume that  $q_*$  minimizes the value  $\|q_* - p\|_\infty^-$  among all such vectors. Since

$$\|q_* - p\|_\infty^+ + \|q_* - p\|_\infty^- = \mu(p) < \|p_* - p\|_\infty^+ + \|p_* - p\|_\infty^- = \|p_* - p\|_\infty,$$

we have  $p_* \neq q_*$  and  $\|q_* - p\|_\infty^- > 0$ . Hence,  $q_*(j) < p(j)$  holds for some  $j \in N$ .

By the translation submodularity (2.4) with  $\alpha = \max_{i \in N} \{p_*(i) - q_*(i)\} - 1$ , we have

$$g(p_*) + g(q_*) \geq g(p_* \wedge (q_* + \alpha \mathbf{1})) + g((p_* - \alpha \mathbf{1}) \vee q_*) = g(p_* - e_Y) + g(q_* + e_Y), \quad (\text{A.2})$$

where  $Y = \arg \max_{i \in N} \{p_*(i) - q_*(i)\}$ . Note that  $\alpha \geq 0$  holds since  $p_*(j) \geq p(j) > q_*(j)$ . Since  $p_*, q_* \in \arg \min g$ , the inequality (A.2) implies  $p_* - e_Y, q_* + e_Y \in \arg \min g$ . We have

$$\|(q_* + e_Y) - p\|_\infty^+ \leq \|q_* - p\|_\infty^+ + \|e_Y\|_\infty^+ = \|q_* - p\|_\infty^+ + 1.$$

As shown below, we have  $X \subseteq Y$  for  $X = \arg \max_{i \in N} \{p(i) - q_*(i)\}$ , implying that

$$\|(q_* + e_Y) - p\|_\infty^- = \|q_* - p\|_\infty^- - 1. \quad (\text{A.3})$$

Hence, we have

$$\|(q_* + e_Y) - p\|_\infty^+ + \|(q_* + e_Y) - p\|_\infty^- \leq \|q_* - p\|_\infty^+ + \|q_* - p\|_\infty^- = \mu(p),$$

which, together with the definition of  $\mu(p)$ , implies  $\|(q_* + e_Y) - p\|_\infty^+ + \|(q_* + e_Y) - p\|_\infty^- = \mu(p)$ . This, however, contradicts the choice of  $q_*$  since (A.3) holds. Therefore, the inequality in (A.1) holds with equality.

To conclude the proof, we show  $X \subseteq Y$ . Since  $p_* - e_Y \in \arg \min g$ , the choice of  $p_*$  implies that  $p_* - e_Y \not\geq p$ . Therefore, it holds that  $p_*(h) = p(h)$  for some  $h \in Y$ . For each  $i \in X$ , we have

$$p_*(i) - q_*(i) \geq p(i) - q_*(i) \geq p(h) - q_*(h) = p_*(h) - q_*(h),$$

where the first inequality is by  $p_* \geq p$  and the second by the definition of  $X$ . The inclusion  $X \subseteq Y$  follows from the inequality above,  $h \in Y$ , and the definition of  $Y$ .

## References

- [1] R.K. Ahuja, T.L. Magnanti, and J.B. Orlin: *Network Flows—Theory, Algorithms and Applications* (Prentice-Hall, Englewood Cliffs, 1993).
- [2] T. Andersson and A. Erlanson: Multi-item Vickrey–English–Dutch auctions. *Games and Economic Behavior*, **81** (2013), 116–129.
- [3] L.M. Ausubel: An efficient dynamic auction for heterogeneous commodities. *American Economic Review*, **96** (2006), 602–629.
- [4] J.M. Bioucas-Dias and G. Valadão: Phase unwrapping via graph cuts. *IEEE Transactions on Image Processing*, **16** (2007), 698–709.
- [5] L. Blumrosen and N. Nisan: Combinatorial auction. In N. Nisan, T. Roughgarden, É. Tardos, and V.V. Vazirani (eds.): *Algorithmic Game Theory* (Cambridge University Press, Cambridge, 2007), 267–299.
- [6] Y.B. Boykov, O. Veksler, and R. Zabih: Fast approximate energy minimization via graph cuts. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **23** (2001), 1222–1239.
- [7] N.-k. Chung and D.-w. Tcha: A dual algorithm for submodular flow problems. *Operations Research Letters*, **10** (1991), 489–495.
- [8] P. Cramton, Y. Shoham, and R. Steinberg: *Combinatorial Auctions* (MIT Press, Cambridge, MA, 2006).
- [9] J. Darbon and M. Sigelle: Image restoration with discrete constrained total variation, part I: Fast and exact optimization. *Journal of Mathematical Imaging and Vision*, **26** (2006), 261–276.
- [10] G. Demange, D. Gale, and M. Sotomayor: Multi-item auctions. *Journal of Political Economy*, **94** (1986), 863–872.
- [11] A.W.M. Dress and W. Wenzel: Valuated matroid: A new look at the greedy algorithm. *Applied Mathematics Letters*, **3** (1990), 33–35.
- [12] A.W.M. Dress and W. Wenzel: Valuated matroids. *Advances in Mathematics*, **93** (1992), 214–250.
- [13] M. Drexler and A. Kleiner: An efficient dynamic auction for general economies with indivisibilities. Working Paper (2015).

- [14] S. Fujishige: *Submodular Functions and Optimization, 2nd edition* (Elsevier, Amsterdam, 2005).
- [15] S. Fujishige and K. Murota: Notes on L-/M-convex functions and the separation theorems. *Mathematical Programming*, **88** (2000), 129–146.
- [16] S. Fujishige, K. Murota, and A. Shioura: Monotonicity in steepest ascent algorithms for polyhedral L-concave functions. *Journal of Operations Research Society of Japan*, **58** (2015), 184–208.
- [17] S. Fujishige and Z. Yang: A note on Kelso and Crawford’s gross substitutes condition. *Mathematics of Operations Research*, **28** (2003), 463–469.
- [18] S. Geman and D. Geman: Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **6** (1984), 721–741.
- [19] F. Gul and E. Stacchetti: Walrasian equilibrium with gross substitutes. *Journal of Economic Theory*, **87** (1999), 95–124.
- [20] F. Gul and E. Stacchetti: The English auction with differentiated commodities. *Journal of Economic Theory*, **92** (2000), 66–95.
- [21] R. Hassin: The minimum cost flow problem: A unifying approach to dual algorithms and a new tree-search algorithm. *Mathematical Programming*, **25** (1983), 228–239.
- [22] H. Hirai: L-extendable functions and a proximity scaling algorithm for minimum cost multiflow problem. *Discrete Optimization*, **18** (2015), 1–37.
- [23] H. Hirai: Discrete convexity and polynomial solvability in minimum 0-extension problems. *Mathematical Programming*, **155** (2016), 1–55.
- [24] W.T. Huh and G. Janakiraman: On the optimal policy structure in serial inventory systems with lost sales. *Operations Research*, **58** (2010), 486–491.
- [25] S. Iwata, L. Fleischer, and S. Fujishige: A combinatorial, strongly polynomial-time algorithm for minimizing submodular functions. *Journal of the ACM*, **48** (2001), 761–777.
- [26] A.S. Kelso and V.P. Crawford: Job matching, coalition formation and gross substitutes. *Econometrica*, **50** (1982), 1483–1504.
- [27] V. Kolmogorov: Primal-dual algorithm for convex markov random fields. Technical Report MSR-TR-2005-117, Microsoft (2005).
- [28] V. Kolmogorov and A. Shioura: New algorithms for convex cost tension problem with application to computer vision. *Discrete Optimization*, **6** (2009), 378–393.
- [29] D. Mishra and D.C. Parkes: Multi-item Vickrey–Dutch auctions. *Games and Economic Behavior*, **66** (2009), 326–347.
- [30] K. Murota: Convexity and Steinitz’s exchange property. *Advances in Mathematics*, **124** (1996), 272–311.
- [31] K. Murota: Discrete convex analysis. *Mathematical Programming*, **83** (1998), 313–371.
- [32] K. Murota: *Discrete Convex Analysis* (Kyoritsu Shuppan, Tokyo, 2001) (in Japanese).
- [33] K. Murota: *Discrete Convex Analysis* (Society for Industrial and Applied Mathematics, Philadelphia, PA, 2003).
- [34] K. Murota: On steepest descent algorithms for discrete convex functions. *SIAM Journal on Optimization*, **14** (2003), 699–707.
- [35] K. Murota: M-convex functions on jump systems: A general framework for minsquare graph factor problem. *SIAM Journal on Discrete Mathematics*, **20** (2006), 213–226.

- [36] K. Murota: *Primer of Discrete Convex Analysis* (Kyoritsu Shuppan, Tokyo, 2007) (in Japanese).
- [37] K. Murota: Recent developments in discrete convex analysis. In W. Cook, L. Lovász and J. Vygen (eds.): *Research Trends in Combinatorial Optimization* (Springer, Berlin, 2009), Chapter 11, 219–260.
- [38] K. Murota and A. Shioura: M-convex function on generalized polymatroid. *Mathematics of Operations Research*, **24** (1999), 95–105.
- [39] K. Murota and A. Shioura: Extension of M-convexity and L-convexity to polyhedral convex functions. *Advances in Applied Mathematics*, **25** (2000), 352–427.
- [40] K. Murota and A. Shioura: Conjugacy relationship between M-convex and L-convex functions in continuous variables. *Mathematical Programming*, **101** (2004), 415–433.
- [41] K. Murota and A. Shioura: Quadratic M-convex and L-convex functions. *Advances in Applied Mathematics*, **33** (2004), 318–341.
- [42] K. Murota and A. Shioura: *Discrete Convex Analysis and Optimization Algorithms* (Asakura Shoten, Tokyo, 2013) (in Japanese).
- [43] K. Murota and A. Shioura: Dijkstra’s algorithm and L-concave function maximization. *Mathematical Programming*, **145** (2014), 163–177.
- [44] K. Murota and A. Shioura: Exact bounds for steepest descent algorithms of L-convex function minimization. *Operations Research Letters*, **42** (2014), 361–366.
- [45] K. Murota, A. Shioura, and Z. Yang: Computing a Walrasian equilibrium in iterative auctions with multiple differentiated items. Extended abstract version in: L. Cai, S.-W. Cheng, and T.W. Lam (eds.): *Proceedings of the 24th International Symposium on Algorithms and Computation (ISAAC 2013)*, LNCS **8283**, 468–478; Full paper version in: Technical Report METR 2013-10, University of Tokyo (2013).
- [46] K. Murota, A. Shioura, and Z. Yang: Time bounds for iterative auctions: A unified approach by discrete convex analysis. *Discrete Optimization*, **19** (2016), 36–62.
- [47] R.T. Rockafellar: *Convex Analysis* (Princeton University Press, Princeton, NJ, 1970).
- [48] H. Saito: The combinatorial English auction of Gul and Stacchetti viewed from discrete convex analysis. Master’s Thesis, Osaka University (2003).
- [49] A. Schrijver: A combinatorial algorithm minimizing submodular functions in strongly polynomial time. *Journal of Combinatorial Theory B*, **80** (2000), 346–355.
- [50] A. Schrijver: *Combinatorial Optimization—Polyhedra and Efficiency* (Springer, Berlin, 2003).
- [51] A. Shioura: L-convex function minimization algorithms: Connection between discrete convex analysis and other fields of research. In A. Tamura (ed.): *Proceedings of 27th RAMP Symposium* (Operations Research Society of Japan, Tokyo, 2015), 17–46 (in Japanese).
- [52] É. Tardos: Generalized matroids and supermodular colourings. In A. Recski and L. Lovász (eds.): *Matroid Theory, Colloquia Mathematica Societatis János Bolyai*, **40** (North-Holland, Amsterdam, 1985), 359–382.
- [53] P. Zipkin: On the structure of lost-sales inventory models. *Operations Research*, **56** (2008), 937–944.
- [54] A. Zomet, A. Levin, S. Peleg, and Y. Weiss: Seamless image stitching by minimizing false edges. *IEEE Transactions on Image Processing*, **15** (2006), 969–977.

Akiyoshi Shioura  
Department of Industrial Engineering and  
Economics  
Tokyo Institute of Technology  
Oh-okayama 2-12-1, Meguro-ku  
Tokyo 152-8500, Japan  
E-mail: [shioura.a.aa@m.titech.ac.jp](mailto:shioura.a.aa@m.titech.ac.jp)